



# JSM Dentistry

---

## JSM Dentistry

---

# JSM Dentistry

[Download Brochure](#)



## Editorial Board



[Stephanos Kyrkanides,](#)

Stony Brook University, USA

[Vaishali Singhal,](#)

University of Medicine and Dentistry of New Jersey, USA

[Jae Hyun Park,](#)

Arizona School of Dentistry & Oral Health, USA

[Xiu-Ping Wang,](#)

Harvard School of Dental Medicine, USA

[Ronald Stephen Brown,](#)

Howard University, USA

[Wenjian Zhang,](#)

University of Texas School of Dentistry at Houston, USA

[Yusuke Shiozawa,](#)

University of Michigan, USA

[Richard L. Jacobson,](#)

FCLA, USA

[Liang Chen,](#)

Bisco Inc., USA

[Catherine Hong Hsu Ling,](#)

University of Michigan, USA

[Vandana Kumar,](#)

University of Missouri-Kansas City, USA

[Rico D. Short,](#)

Medical College of Georgia School of Dentistry, USA

[Ying Liu,](#)

University of Missouri at Kansas City, USA

[Andrei Barasch,](#)

Winthrop University Hospital, USA

[Shuying \(Sheri\) Yang,](#)

University at Buffalo, USA

[Luiz Andre Pimenta,](#)

University of North Carolina, USA



[Rohitashw Kumar,](#)

State University of New York at Buffalo, USA

[Maobin Yang,](#)

Temple University, USA

[Cemil Yesilsoy,](#)

Temple University, USA

[Michael J. Metz,](#)

University of Louisville School of Dentistry, USA

[Rajendra Prasad Settem,](#)

State University of New York at Buffalo, USA

[Timothy T. Wheeler,](#)

University of Florida Gainesville, USA

[Peter T. Gakunga,](#)

University of Texas Health Science Center San Antonio, USA

[Chung H. Kau,](#)

University of Alabama, USA

[Robert Michael Block,](#)

Meharry Medical College, USA

[Harold O. Jackson,](#)

Meharry Medical College, USA

[Ping Zhang,](#)

University of Alabama at Birmingham, USA

[Laurence Shapiro,](#)

Nova Southeastern University, USA

[Karen Kulikowski,](#)

The State University of New Jersey, USA

[Maria A. Hernandez,](#)

Nova Southeastern University, USA

[Bing-Yan Wang,](#)

University of Texas Health Science Center at Houston, USA

[LaQuia A. Walker,](#)

Indiana University School of Dentistry, USA



[University of Manitoba, USA](#)

[Priyanshi Ritwik,](#)

LSU School of Dentistry, USA

[David E. Simmons,](#)

LSU Health Sciences Center, USA

[Peter M. Gershenson,](#)

Albert Einstein College of Medicine, USA

[Charles Lekic,](#)

University Of Manitoba, USA

[Laurence James Walsh,](#)

The University of Queensland, Australia

[Anthony L. Maganzini,](#)

Albert Einstein College of Medicine, USA

[Zheng Xu,](#)

University of Texas Health Science Center, USA

[Sorin Uram-Tuculescu,](#)

VCU School of Dentistry, USA

[Varkha Agrawal,](#)

NorthShore University Healthsystem, USA

[Tan Kai Soo,](#)

National University of Singapore, Singapore

[Hamid Nurrohman,](#)

University of California, USA

[Masataka Sunohara,](#)

The Nippon Dental University, Japan

[Md. Sofiqul Islam,](#)

Tokyo Medical & Dental University, Japan

[Tada-aki Kudo,](#)

Tohoku University Graduate School of Dentistry, Japan

[Harry Huiz Peeters,](#)

Laser Research Center Bandung, Indonesia



[Tang Thean Hock,](#)

[Universiti Sains Malaysia, Malaysia](#)

[Abhishek Parolia,](#)

International Medical University, Malaysia

[Yukinori Maruo,](#)

Okayama University, Japan

[Prasansuttiporn Taweesak,](#)

Chiang Mai University, Thailand

[Junichi Asaumi,](#)

Okayama University Graduate School of Medicine, Japan

[Akiko Kato,](#)

Aichi-Gakuin University, Japan

[Jun Murakami,](#)

Okayama University Graduate School of Medicine, Japan

[Theodore Eliades,](#)

University of Zurich, Switzerland

[Ronald Wigler,](#)

Rambam Health Care Campus, Israel

[Azin Parsa,](#)

Academic centre for dentistry Amsterdam, The Netherlands

[Ying-Chu Lin,](#)

Kaohsiung Medical University, Taiwan

[Helder Nunes Costa,](#)

Kanagawa Dental University, Japan

[K Kamburoglu,](#)

Ankara University, Turkey

[Vajendra Joshi,](#)

Showa University, Japan

[Aysegu Olmez,](#)

University of Gazi, Turkey

[Kin-ya Kubo,](#)

Seijoh University Graduate School of Health Care Studies, Japan



[Naoki Kodama,](#)

Universite de Montreal, Canada

[Akito Tomomura,](#)

Meikai University School of Dentistry, Japan

## Editor Comment Section

Nihon University School of Dentistry, Japan

Full Name

Email Id

Pravara Institute of Medical Sciences, India

Your Comment

F

[Plauto Christopher Aranha Watanabe,](#)

University of Sao Paulo, Brazil

Submit

## Guidelines

» [Editor](#)

» [Reviewer](#)

Mersin University School of Dentistry, Turkey

If you are an experienced academic with an interest in open Science please.. share your details.....

[Aysa Ayali,](#)

Near East University, Turkey

Join with us

[Mei-Chi Chang,](#)

Chang Gung University of Science and Technology, Taiwan

## Editor Articles

📅 September 20, 2020



[Download Brochure](#)[f](#) [t](#) [G+](#) [in](#) [p](#)

## 2015 Table of Contents

### Volume 3 - Issue 1 Articles

#### **[Fracture Strength of Flared Roots Restored with an Innovative Glass Fiber Post Technique](#)**

Article

[Andréa de Azevedo Brito Conceição](#), [Armiliana Soares Nascimento](#), [Ewerton Nocchi Conceição](#), [Fernanda Maria Barros Guerra](#), [José Eurípedes De Oliveira](#), [Rodivan Braz](#)

September 14, 2015 | Pages : 205 - 450

**Abstract** Proposition: evaluate the fracture strength of endodontically treated flared roots after different techniques of intraradicular restoration.  
Methods: Thirty human upper canines with similar root dimensions were sectioned transversely and the root....

[Abstract](#) | [EPub](#) | [XML](#) | [Full text](#) | [PDF](#)[f](#) [t](#) [G+](#) [in](#)

---

#### **[Sealing Capacity of Different Materials used as Cervical Barrier during Internal Walking Bleaching Technique](#)**

Article

[José Renato Cavalcanti de Queiroz](#), [Juliana Fernandes de Moraes](#), [Cláudia Tavares Machado](#), [Leticia Virgínia de Freitas Chaves](#), [Lorena Martins de Sá Torres](#), [Marinna Ramalho Rocha de Melo](#), [Vanessa Suzana de Freitas Chaves](#)

August 06, 2015 | Pages : 205 - 450

**Abstract** Objectives: This study evaluated the sealing ability of different temporary restorative materials used as a barrier in the passage of oxygen generated during internal bleaching technique mediated by different time intervals. Material and methods.....

[Abstract](#) | [EPub](#) | [XML](#) | [Full text](#) | [PDF](#)[f](#) [t](#) [G+](#) [in](#)

---

#### **[Goldenhar Syndrome – A Literature Review Shruti Sinha1 \\*, Ashish Kumar Singh2 , A](#)**

Article

[Anshul Mehra](#), [Ashish Kumar Singh](#), [Rahul Singh](#), [Shruti Sinha](#)

July 06, 2015 | Pages : 205 - 450

**Abstract** Golden har syndrome is a hereditary condition which is characterized by preauricular appendages, fistulas, and epibulbar dermoids. It not only involves the facial structures, but also includes renal, genitourinary, cardiac, and skeletal anomalies.....

[Abstract](#) | [EPub](#) | [XML](#) | [Full text](#) | [PDF](#)[f](#) [t](#) [G+](#) [in](#)

---

#### **[Evaluation of Periodontal Parameters in Patients Peritoneal Dialysis, Hemodialysis and Renal Transplantation](#)**

Article

^



[Mustafa Kele?](#), [Hülya Aksoy](#), [Gülnihal Emrem Doğan](#)

July 04, 2015 | Pages : 205 - 450

**Abstract** Objectives: Chronic renal failure (CRF) is defined as a progressive decline in renal function associated with a reduced glomerular filtration rate. Treatment of this failure consists of hemodialysis (HD), peritoneal dialysis (PD) and renal transp.....

[Abstract](#) | [EPub](#) | [XML](#) | [Full text](#) | [PDE](#)

[f](#) [t](#) [G+](#) [in](#)

### [Association between Tooth Loss and the Development of Dementia on the Indonesian Elders](#)

Article

[Adang Bachtiar](#), [Anton Rahardjo](#), [Asyurati Asia](#), [Lindawati Kusdhany](#)

July 04, 2015 | Pages : 205 - 450

**Abstract** Tooth loss may be a modifiable risk factor for dementia, but the causal relationship has not been evaluated sufficiently. The purpose of this study was to investigate the association between tooth loss and dementia on the Indonesian elder. The sub.....

[Abstract](#) | [EPub](#) | [XML](#) | [Full text](#) | [PDE](#)

[f](#) [t](#) [G+](#) [in](#)

### [Incidental Finding of Carotid Artery Calcification in a Prosthodontic Patient](#)

Article

[Bader K Alzarea](#)

June 16, 2015 | Pages : 205 - 450

**Abstract**

[Abstract](#) | [EPub](#) | [XML](#) | [Full text](#) | [PDE](#)

[f](#) [t](#) [G+](#) [in](#)

### [Palatal Swelling Secondary to Radicular Cyst](#)

Article

[Santosh R Patil](#)

January 05, 2015 | Pages : 205 - 450

**Abstract**

[Abstract](#) | [EPub](#) | [XML](#) | [Full text](#) | [PDE](#)

[f](#) [t](#) [G+](#) [in](#)

## Browse by Year

2015



See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/291355704>

# Association between Tooth Loss and the Development of Dementia on the Indonesian Elders

Article · July 2015

CITATIONS

2

READS

207

4 authors, including:



**Lindawati Kusdhany**

University of Indonesia

64 PUBLICATIONS 393 CITATIONS

[SEE PROFILE](#)



**Anton Rahardjo**

University of Indonesia

92 PUBLICATIONS 612 CITATIONS

[SEE PROFILE](#)



**Adang Bachtiar**

University of Indonesia

73 PUBLICATIONS 186 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Tooth Eruption Systematic Review [View project](#)



Niigata Elderly Study [View project](#)

## Research Article

# Association between Tooth Loss and the Development of Dementia on the Indonesian Elders

Asyurati Asia \*, Lindawati Kusdhany , Anton Rahardjo and Adang Bachtiar

<sup>1</sup>Department of Dentistry, Universitas Indonesia, Indonesia

<sup>2</sup>Department of Public Health, Universitas Indonesia, Indonesia

## Abstract

Tooth loss may be a modifiable risk factor for dementia, but the causal relationship has not been evaluated sufficiently. The purpose of this study was to investigate the association between tooth loss and dementia on the Indonesian elder. The subjects consisted of 208 volunteers (36 males and 172 females) aged 60 years or older. The mini mental state examination (MMSE; score range: 0-30) is used as a screening test for cognitive impairment. In the logistic regression analysis, odds ratio (OR) of cognitive impairment (MMSE score < 23) were 6.34 (95% CI, 2.76 to 14.55) and 2.82 (95% CI, 1.30 to 6.10) for subjects who had remaining < 10 teeth and those who had remaining 11-19 teeth, respectively, when compared with subjects who had remaining > 20 teeth. It can be concluded that tooth loss have association with dementia in Indonesian subjects.

## ABBREVIATIONS

MMSE: Mini Mental State Examination

## INTRODUCTION

Dementia is a syndrome due to disease of the brain, usually of a chronic or progressive nature, in which there is disturbance of multiple higher cortical functions, including memory, thinking, orientation, comprehension, calculation, learning capacity, language, and judgment. In this syndrome, the impairments of cognitive function are commonly accompanied, and occasionally preceded, by deterioration in emotional control, social behavior, or motivation. Moreover, this syndrome occurs in a large number of conditions primarily or secondarily affecting the brain [1].

Numerous studies have linked dementia to the subsequent deterioration of oral health and the most significant public health problem in ageing community. It is estimated that there are 24.3 million people with dementia worldwide and projected there are 4.6 million new cases raising each year. By the year 2040, it is expected that dementia will influence more than 81 million people worldwide [2]. In the year 2010, the elderly population whose age is more than 60 years in Indonesia is predicted at approximately 19 million, while the actual prevalence of dementia is not known. On the other hand, according to the Alzheimer's disease International projection, the number of dementia in Indonesia was estimated to be 606,100 people in 2005 with

## \*Corresponding author

Asyurati Asia, Department of Dentistry, Universitas Indonesia, Jakarta, Indonesia, Tel: 62-818-479-415; Email: asyuratiasia@gmail.com

**Submitted:** 09 December 2014

**Accepted:** 02 July 2015

**Published:** 04 July 2015

**ISSN:** 2333-7133

## Copyright

© 2015 Asia et al.

OPEN ACCESS

## Keywords

- Tooth loss
- Dementia
- Elderly

annual incidence cases of 191,400. This number was projected to be 1,016,800, with annual new cases of 314,000 by the year 2020 [3,4]. Several studies have reported that tooth loss is may be associated with dementia in elders [5-8]. To our knowledge, however, only a small number of community-based studies have investigated the causal relationships between tooth loss and dementia with cognitive impairment. Further understanding of the modifiable risk factors of dementia will be helpful to reduce the rapid incidence of disease. The purpose of this study was to investigate the association between tooth loss and dementia in an elderly Indonesian population.

## MATERIALS AND METHODS

The subjects consisted of 208 volunteers (36 males and 172 females) aged 60 years or older. Fifty three participants (22 males and 31 females) were recruited from Sukabumi rural area while 155 participants (14 males and 141 females) were recruited from urban area from the city of Jakarta (capital of Indonesia) and Depok a satellite of Jakarta. All of the participants were living in their own homes and most of them were able to walk independently. This study was approved by the Ethics Committee of Faculty of Dentistry, Universitas Indonesia. Written informed consent was obtained from each of the subjects before their participation in the study.

The Mini Mental State Examination (MMSE) is widely used as a global test of cognitive function to screen for dementia and assesses orientation, immediate memory, attention and calculation, short-term recall, and language [9]. Based on the original MMSE, we developed a modified and culturally adapted MMSE according to Indonesian subjects. The MMSE (score range, 0-30) is used as a screening test for cognitive impairment. The Recall test (score range, 0-3) is a sub-item of the IMMSE that evaluates the impairment of recent memory. Subjects were instructed to recall three unrelated objects that they were previously instructed to remember. The MMSE was carried out by two interviewers who were formally trained by a psychiatrist. Cognitive impairment was classified as an MMSE score of less than 24 [9, 10].

A dental examination was carried out by two dentists calibrated regarding the techniques, with the dentist and the subject in a sitting position under artificial lighting. The number of teeth was recorded for each subject. The remaining teeth were defined as healthy, carious, or treated teeth, inclusive of completely erupted third molars. Severe carious teeth and very loose teeth that were indicated for extraction were excluded as remaining teeth. We categorized the remaining teeth as twenty teeth or more, eleven to nineteen, ten teeth or less.

We classified that having the status of hypertension when a participant's having systolic blood pressure of 140 mm Hg or greater and/or diastolic blood pressure of 90 mm Hg or greater. We categorized participant's non-fasting glucose level as 200 mg/dL or over and below than 200 mg/dL.

Statistical analyses were performed using Confidence Interval Analysis Software for Windows (ver. 2.0, Trevor Bryant, University of Southampton). Logistic regression analysis was carried out with low MMSE score as dependent variables and other variables were used as independent variables. The number of remaining teeth was used as continuous variables in one model and categorical variables in the other model.

## RESULTS AND DISCUSSION

### Results

All of the 208 participants aged 60 years old and above (36 men and 172 women) have completed all the examination processes. The subjects were divided into two groups based on their MMSE scores. The demographic characteristics of subjects, the socioeconomic characteristics and the number of remaining teeth are shown in Table 1.

The mean age (SD) of male and female subjects were 74.4 (8.5) and 66.8 (6.9) years old, respectively. The mean MMSE score (SD) of subjects with low MMSE score ( $\leq 23$ ) and high MMSE score ( $\geq 24$ ) were 20.1 (8.8) and 27.0 (2.0), respectively. The proportions of men and women having low MMSE were found 25% (9/36) and 25.6% (44/172), respectively. Subjects living in the rural area were more likely to have a lower MMSE score. The proportions of urban and rural elders having low MMSE were found 20.7% (32/156) and 39.6% (21/53) respectively. Most of subjects with low MMSE were having 6 years or lower education experience (37.8%). The proportion of subjects with low MMSE among 7 - 12 years and  $\geq 13$  years educational experience groups were 19.4% (21/108) and 5.6% (1/18), respectively. Monthly allowances among subjects with low MMSE was found as low as IDR 512,903 while in the high MMSE group was found as high as IDR 922,288.

The mean (SD) number of remaining teeth in subjects with low MMSE and high MMSE were 14.65 (7.96) and 21.5 (8.02), respectively. In the logistic regression analysis, odds ratio (OR) of cognitive impairment (MMSE score  $\leq 23$ ) were 6.34 (95% CI, 2.76 to 14.55) and 2.82 (95% CI, 1.30 to 6.10) for subjects who had remaining  $\leq 10$  teeth and those who had remaining 11 - 19 teeth, respectively, when compared with subjects who had remaining  $\geq 20$  teeth.

### Discussion

The result of this community-based survey revealed that the

**Table 1:** Distribution of demographic, socioeconomic, and number of remaining teeth status.

Parameter	Low MMSE ( $\leq 23$ )		High MMSE ( $\geq 24$ )	
	Mean	(SD)	Mean	(SD)
Age (years), mean (SD)	73.45	(8.92)	66.29	(5.38)
Sex, number (%)				
males	9	(25.0)	27	(75.0)
females	44	(25.6)	128	(74.4)
Region, number (%)				
urban	32	(15.38)	123	(59.13)
rural	21	(10.10)	32	(15.38)
Length of education, number (%)				
$\geq 13$ yrs	1	(5.6)	17	(94.4)
7 - 12 yrs	21	(19.4)	87	(80.6)
$\leq 6$ yrs	31	(37.8)	51	(62.2)
Allowances (IDR/month), mean (SD)	512,903	(507,452)	1,541,389	(922,288)
Number of remaining teeth, mean (SD)	14.65	(7.96)	21.50	(8.02)

**Abbreviations:** IDR: Indonesian Rupiah; MMSE: Mini Mental State Examination; SD: Standard Deviation

prevalence of a low MMSE score was significantly associated with low number of remaining teeth. These results are consistent with those of previous studies demonstrating that tooth loss was associated with decreased cognitive function [5,7,8]. Many factors may contribute to cognitive impairment in elders, including cardiovascular disease, education and social economy status [10].

The other main findings of this study are that a low level of education and a low income were significantly associated with low MMSE score and increased risk of dementia. Our results are in agreement with previous findings in a study of education and dementia in a Japanese population[7]. Different mechanisms have been suggested to explain this association. Katzman [11] proposed that education might increase brain reserves by increasing synaptic density in the neocortical association cortex. Stern et al. extended the cognitive reserve hypothesis and took into account the possible beneficial influence of mental activity throughout the entire life span, where occupational attainment along with level of education could influence the risk of dementia's disease [12]. It is known that education is an indicator of socioeconomic status. This socioeconomic status acting over the life course have been found to affect health and risk of cardiovascular disease. Cumulative economic hardship has been found to lead to poorer cognitive functioning and dementia.

The association between loss of molars and spatial memory deficits was first described in an animal study using senile rats [13]. In that study, after molar extraction, rats were maintained with diet powder and were subjected to behavioral testing to evaluate their spatial memory. Animals with extracted molar teeth showed worse behavioral performance compared to control animals maintained with solid diet. Mastication is the main function of stomatognathic system and the relationship between mastication and the central nervous system (CNS), especially the cognitive area, is still the subject of several studies. Numerous studies have revealed an association between mastication and cognitive functions [10, 13-16]. The possible associations between masticatory hypofunction and cognitive insufficiency have been demonstrated in several animal studies [16-19]. Rats in those studies were induced masticatory hypo function by performing extraction or occlusal reduction of the molar teeth rather than a long-term soft diet [16,17,19]. The occlusal hypo function caused degenerative changes in their periodontal mechanoreceptors [20] suggesting a suppression of sensory stimulation from the periodontal ligament during chewing. In those experiments, animals showed poor performance on memory and learning evaluation.

Histologically, the hippocampus, the main CNS region, that affected by masticatory hypofunction was reported by several studies [16,21]. The hippocampus is included in some kinds of memory, especially which relates to details and precise events [16, 21]. Recently, an animal study using senescence-accelerated mouse prone 8 (SAMP8) examined whether reduced mastication from a young age mice affects hippocampal-dependent cognitive function[22] (Table 2). This study found that loss of masticatory function in early life causes malnutrition and chronic stress and impairs the ability to recognize novel objects. Hyper activation and lateralized rotational behavior are commonly observed with dysfunction of the dopaminergic system, therefore,

**Table 2:** Logistic regression model for predicting lower MMSE scores (≤23).

Variables	OR	95% CI	P value
<b>Age</b>			
≤ 65 years (reference)	1.00		
≥ 66 years	8.86	3.76 to 20.87	< 0.05
<b>Sex</b>			
males (reference)	1.00		
females	1.03	0.45 to 2.36	> 0.05
<b>Region</b>			
Urban (reference)	1.00		
Rural	2.52	1.29 to 4.95	< 0.05
<b>Length of education</b>			
13 years or over (reference)	1.00		
7 - 12 years	4.10	0.52 to 32.60	> 0.05
6 years or under	10.33	1.31 to 81.53	< 0.05
<b>Allowances (IDR/month)</b>			
≥ 1,500,000 (reference)	1.00		
< 1,500,000	3.01	1.56 to 5.82	< 0.05
<b>Blood pressure</b>			
Normotension (reference)	1.00		
Hypertension	1.38	0.68 to 2.81	< 0.05
<b>Non fasting glucose level (mg/dL)</b>			
< 200 (reference)	1.00		
≥ 200	0.17	0.02 to 1.29	> 0.05
<b>Number of remaining teeth</b>			
≥ 20 (reference)	1.00		
11-19	2.82	1.30 to 6.10	< 0.05
≤ 10	6.34	2.76 to 14.55	< 0.05

**Abbreviations:** CI: Confidence Interval; IDR: Indonesian Rupiah; Mg/dL: Milligram per Deciliter; OR: Odds Ratio.

reduced masticatory function may deplete the mesolimbic and mesocorticolimbic dopaminergic systems to impair the cognitive functions of selective attention and recognition memory in the prefrontal cortex and the hippocampus [22-24].

Masticatory dysfunction also leads to various morphologic changes in the hippocampus and cerebral cortex, e.g., the molar less condition and soft-diet feeding decrease the brain cells including synapses in the hippocampus and parietal cortex [25], as well as neurotrophic receptor expression in hippocampal CA1 and CA3 regions[26]. Hypertrophy and hyperplasia of glial fibrillary acid protein-labeled astrocytes in the CA1 region [27, 28] and suppressed cell proliferation in the hippocampal dentate gyrus [29,30]also occur in mice with learning deficits. These behavioral and hippocampal morphologic changes are similar to the changes associated with advanced age [31,32] and masticatory dysfunction may therefore accelerate the aging process in the hippocampus.

Neuroimaging clinical trials using functional MRI and positron emission tomography indicate that several regions of the brain

are activated during mastication, including the cortex, thalamus, striatum, and cerebellum [33-35]. Functional MRI evaluation on neuronal activity in the brain indicated that chewing activity increases the blood oxygen level-dependent signals in the right premotor cortex, precuneus, thalamus, hippocampus and inferior parietal lobe [36]. Furthermore, chewing activity was reported significantly enhances memory in aged subjects [37]. Cerebellar functions also seem to be influenced by masticatory activity [34]. Recently, Quintero and others [38] have reported that the functional connections between cerebellum and cortical areas during masticatory activity [36-38]. It has been well documented that the cerebellum is responsible for motor planning, including the force needed to perform motor tasks. However, it is not clear yet what is the mechanism of reduced input activity due to masticatory dysfunction differ from the effects of a reduction in other types of sensory stimuli in the aging hippocampus. To elucidate the mechanisms underlying the specific effects of masticatory dysfunction in aging hippocampus, it will be necessary to compare the impact of mastication and environmental enrichment or physical exercise on hippocampal function.

## CONCLUSION

In summary, there are significant relationships were found between the number of remaining teeth and cognitive function. Edentulism may be predictors of dementia late in life.

## ACKNOWLEDGEMENTS

This work was supported by Universitas Trisakti, Jakarta Indonesia. The authors declare no potential conflicts of interest with respect to the authorship and/or publication of this article. We thank Dr. Widijanto Sudhana, M. Kes, Faculty of Dentistry, Universitas Trisakti for constructive statistical comments which have assisted us in preparing this article.

## REFERENCES

1. WHO. Dementia: a public health priority. Geneva: World Health Organization; 2012.
2. Ferri CP, Prince M, Brayne C, Brodaty H, Fratiglioni L, Ganguli M, et al. Global prevalence of dementia: a Delphi consensus study. *Lancet*. 2005; 366: 2112-2117.
3. International Alzheimer's Disease. The Prevalence of Dementia Worldwide. 2008.
4. Stein PS, Desrosiers M, Donegan SJ, Yepes JF, Kryscio RJ. Tooth loss, dementia and neuropathology in the Nun study. *J Am Dent Assoc*. 2007; 138: 1314-1322.
5. Okamoto N, Morikawa M, Okamoto K, Habu N, Iwamoto J, Tomioka K, et al. Relationship of tooth loss to mild memory impairment and cognitive impairment: findings from the Fujiwara-kyo study. *Behav Brain Funct*. 2010; 6: 77.
6. Kaye EK, Valencia A, Baba N, Spiro A 3rd, Dietrich T, Garcia RI. Tooth loss and periodontal disease predict poor cognitive function in older men. *J Am Geriatr Soc*. 2010; 58: 713-718.
7. Saito Y, Sugawara N, Yasui-Furukori N, Takahashi I, Nakaji S, Kimura H. Cognitive function and number of teeth in a community-dwelling population in Japan. *Ann Gen Psychiatry*. 2013; 12: 20.
8. Park H, Suk SH, Cheong JS, Lee HS, Chang H, Do SY, Kang JS. Tooth loss may predict poor cognitive function in community-dwelling adults

- without dementia or stroke: the PRESENT project. *J Korean Med Sci*. 2013; 28: 1518-1521.
9. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975; 12: 189-198.
10. Qiu C, De Ronchi D, Fratiglioni L. The epidemiology of the dementias: an update. *Curr Opin Psychiatry*. 2007; 20: 380-385.
11. Katzman R. Education and the prevalence of dementia and Alzheimer's disease. *Neurology*. 1993; 43: 13-20.
12. Stern Y, Gurland B, Tatemichi TK, Tang MX, Wilder D, Mayeux R. Influence of education and occupation on the incidence of Alzheimer's disease. *JAMA*. 1994; 271: 1004-1010.
13. Kato T, Usami T, Noda Y, Hasegawa M, Ueda M, Nabeshima T. The effect of the loss of molar teeth on spatial memory and acetylcholine release from the parietal cortex in aged rats. *Behav Brain Res*. 1997; 83: 239-242.
14. Miura H, Miura K, Mizugai H, Arai Y, Umenai T, Isogai E. Chewing ability and quality of life among the elderly residing in a rural community in Japan. *J Oral Rehabil*. 2000; 27: 731-734.
15. Miura H, Yamasaki K, Kariyasu M, Miura K, Sumi Y. Relationship between cognitive function and mastication in elderly females. *J Oral Rehabil*. 2003; 30: 808-811.
16. Scherder E, Posthuma W, Bakker T, Vuijk PJ, Lobbezoo F. Functional status of masticatory system, executive function and episodic memory in older persons. *J Oral Rehabil*. 2008; 35: 324-336.
17. Ono Y, Yamamoto T, Kubo KY, Onozuka M. Occlusion and brain function: mastication as a prevention of cognitive dysfunction. *J Oral Rehabil*. 2010; 37: 624-640.
18. Kato T, Usami T, Noda Y, Hasegawa M, Ueda M, Nabeshima T. The effect of the loss of molar teeth on spatial memory and acetylcholine release from the parietal cortex in aged rats. *Behav Brain Res*. 1997; 83: 239-242.
19. Kushida S, Kimoto K, Hori N, Toyoda M, Karasawa N, Yamamoto T. Soft-diet feeding decreases dopamine release and impairs aversion learning in Alzheimer model rats. *Neurosci Lett*. 2008; 439: 208-211.
20. Ekuni D, Tomofuji T, Irie K, Azuma T, Endo Y, Kasuyama K, et al. Occlusal disharmony increases amyloid- $\beta^2$  in the rat hippocampus. *Neuromolecular Med*. 2011; 13: 197-203.
21. Yamamoto T, Hirayama A. Effects of soft-diet feeding on synaptic density in the hippocampus and parietal cortex of senescence-accelerated mice. *Brain Res*. 2001; 902: 255-263.
22. Muramoto T, Takano Y, Soma K. Time-related changes in periodontal mechanoreceptors in rat molars after the loss of occlusal stimuli. *Arch Histol Cytol*. 2000; 63: 369-380.
23. Squire LR1, Stark CE, Clark RE. The medial temporal lobe. *Annu Rev Neurosci*. 2004; 27: 279-306.
24. Kawahata M, Ono Y, Ohno A, Kawamoto S, Kimoto K, Onozuka M. Loss of molars early in life develops behavioral lateralization and impairs hippocampus-dependent recognition memory. *BMC Neurosci*. 2014; 15: 4.
25. Yamamoto T, Hirayama A. Effects of soft-diet feeding on synaptic density in the hippocampus and parietal cortex of senescence-accelerated mice. *Brain Res*. 2001; 902: 255-263.
26. Yamazaki K, Wakabayashi N, Kobayashi T, Suzuki T. Effect of tooth loss on spatial memory and trkB-mRNA levels in rats. *Hippocampus*. 2008; 18: 542-547.
27. Onozuka M, Watanabe K, Nagasaki S, Jiang Y, Ozono S, Nishiyama

- K, et al. Impairment of spatial memory and changes in astroglial responsiveness following loss of molar teeth in aged SAMP8 mice. *Behav Brain Res.* 2000; 108: 145-155.
28. Watanabe K, Tonosaki K, Kawase T, Karasawa N, Nagatsu I, Fujita M, et al. Evidence for involvement of dysfunctional teeth in the senile process in the hippocampus of SAMP8 mice. *Exp Gerontol.* 2001; 36: 283-295.
29. Aoki H, Kimoto K, Hori N, Toyoda M. Cell proliferation in the dentate gyrus of rat hippocampus is inhibited by soft diet feeding. *Gerontology.* 2005; 51: 369-374.
30. Kubo KY, Sasaguri K, Ono Y, Yamamoto T, Takahashi T, Watanabe K, et al. Chewing under restraint stress inhibits the stress-induced suppression of cell birth in the dentate gyrus of aged SAMP8 mice. *Neurosci Lett.* 2009; 466: 109-113.
31. Kitraki E, Bozas E, Philippidis H, Stylianopoulou F. Aging-related changes in IGF-II and c-fos gene expression in the rat brain. *Int J Dev Neurosci.* 1993; 11: 1-9.
32. DeToledo-Morrell L GYaMF. Age-dependent alteration in hippocampal synaptic plasticity: relation to memory disorders. *Neurobiol Aging Neurobiol Aging.* 1993; 14: 441-446.
33. Momose T, Nishikawa J, Watanabe T, Sasaki Y, Senda M, Kubota K, et al. Effect of mastication on regional cerebral blood flow in humans examined by positron-emission tomography with  $\mu$ O-labelled water and magnetic resonance imaging. *Arch Oral Biol.* 1997; 42: 57-61.
34. Onozuka M, Fujita M, Watanabe K, Hirano Y, Niwa M, Nishiyama K, et al. Mapping brain region activity during chewing: a functional magnetic resonance imaging study. *J Dent Res.* 2002; 81: 743-746.
35. Onozuka M, Fujita M, Watanabe K, Hirano Y, Niwa M, Nishiyama K, et al. Age-related changes in brain regional activity during chewing: a functional magnetic resonance imaging study. *J Dent Res.* 2003; 82: 657-660.
36. Hirano Y, Obata T, Kashikura K, Nonaka H, Tachibana A, Ikehira H, et al. Effects of chewing in working memory processing. *Neurosci Lett.* 2008; 436: 189-192.
37. Onozuka M HY, Tachibana A, Kim W, Ono Y, Sasaguri K, Kubo K, Niwa M, et al. Interactions between chewing and brain activities in human. *Novel Trends in Brain Science Novel Trends in Brain Science.* 2007; 99-113.
38. Quintero A, Ichesco E, Schutt R, Myers C, Peltier S, Gerstner GE. Functional connectivity of human chewing: an fMRI study. *J Dent Res.* 2013; 92: 272-278.

#### Cite this article

Asia A, Kusdhany L, Rahardjo A, Bachtiar A (2015) Association between Tooth Loss and the Development of Dementia on the Indonesian Elders. *JSM Dent* 3(1): 1050.

# Association between Tooth Loss and the Development of Dementia on the Indonesian Elders

*by Asyurati Asia*

---

**Submission date:** 21-Dec-2022 11:08AM (UTC+0700)

**Submission ID:** 1985334288

**File name:** ToothLossandtheDevelopmentofDementiaontheIndonesianElders\_1.pdf (430.21K)

**Word count:** 3891

**Character count:** 20637



See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/291355704>

# Association between Tooth Loss and the Development of Dementia on the Indonesian Elders

Article · July 2015

CITATIONS

2

READS

207

4 authors, including:



**Lindawati Kusdhary**  
University of Indonesia

64 PUBLICATIONS 393 CITATIONS

[SEE PROFILE](#)



**Anton Rahardjo**  
University of Indonesia

92 PUBLICATIONS 612 CITATIONS

[SEE PROFILE](#)



**Adang Bachtiar**  
University of Indonesia

73 PUBLICATIONS 186 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Tooth Eruption Systematic Review [View project](#)



Niigata Elderly Study [View project](#)

All content following this page was uploaded by [Anton Rahardjo](#) on 22 January 2016.

The user has requested enhancement of the downloaded file.

## Research Article

# Association between Tooth Loss and the Development of Dementia on the Indonesian Elders

Asyurati Asia \*, Lindawati Kusdhany , Anton Rahardjo and Adang Bachtiar

<sup>1</sup>Department of Dentistry, Universitas Indonesia, Indonesia

<sup>2</sup>Department of Public Health, Universitas Indonesia, Indonesia

**\*Corresponding author**

Asyurati Asia, Department of Dentistry, Universitas Indonesia, Jakarta, Indonesia, Tel: 62-818-479-415; Email: asyuratiasia@gmail.com

Submitted: 09 December 2014

Accepted: 02 July 2015

Published: 04 July 2015

ISSN: 2333-7133

Copyright

© 2015 Asia et al.

OPEN ACCESS

**Keywords**

- Tooth loss
- Dementia
- Elderly

**Abstract**

Tooth loss may be a modifiable risk factor for dementia, but the causal relationship has not been evaluated sufficiently. The purpose of this study was to investigate the association between tooth loss and dementia on the Indonesian elder. The subjects consisted of 208 volunteers (36 males and 172 females) aged 60 years or older. The mini mental state examination (MMSE; score range: 0-30) is used as a screening test for cognitive impairment. In the logistic regression analysis, odds ratio (OR) of cognitive impairment (MMSE score < 23) were 6.34 (95% CI, 2.76 to 14.55) and 2.82 (95% CI, 1.30 to 6.10) for subjects who had remaining < 10 teeth and those who had remaining 11-19 teeth, respectively, when compared with subjects who had remaining > 20 teeth. It can be concluded that tooth loss have association with dementia in Indonesian subjects.

**ABBREVIATIONS**

MMSE: Mini Mental State Examination

**INTRODUCTION**

Dementia is a syndrome due to disease of the brain, usually of a chronic or progressive nature, in which there is disturbance of multiple higher cortical functions, including memory, thinking, orientation, comprehension, calculation, learning capacity, language, and judgment. In this syndrome, the impairments of cognitive function are commonly accompanied, and occasionally preceded, by deterioration in emotional control, social behavior, or motivation. Moreover, this syndrome occurs in a large number of conditions primarily or secondarily affecting the brain [1].

Numerous studies have linked dementia to the subsequent deterioration of oral health and the most significant public health problem in ageing community. It is estimated that there are 24.3 million people with dementia worldwide and projected there are 4.6 million new cases raising each year. By the year 2040, it is expected that dementia will influence more than 81 million people worldwide [2]. In the year 2010, the elderly population whose age is more than 60 years in Indonesia is predicted at approximately 19 million, while the actual prevalence of dementia is not known. On the other hand, according to the Alzheimer's disease International projection, the number of dementia in Indonesia was estimated to be 606,100 people in 2005 with

annual incidence cases of 191,400. This number was projected to be 1,016,800, with annual new cases of 314,000 by the year 2020 [3,4]. Several studies have reported that tooth loss is may be associated with dementia in elders [5-8]. To our knowledge, however, only a small number of community-based studies have investigated the causal relationships between tooth loss and dementia with cognitive impairment. Further understanding of the modifiable risk factors of dementia will be helpful to reduce the rapid incidence of disease. The purpose of this study was to investigate the association between tooth loss and dementia in an elderly Indonesian population.

**MATERIALS AND METHODS**

The subjects consisted of 208 volunteers (36 males and 172 females) aged 60 years or older. Fifty three participants (22 males and 31 females) were recruited from Sukabumi rural area while 155 participants (14 males and 141 females) were recruited from urban area from the city of Jakarta (capital of Indonesia) and Depok a satellite of Jakarta. All of the participants were living in their own homes and most of them were able to walk independently. This study was approved by the Ethics Committee of Faculty of Dentistry, Universitas Indonesia. Written informed consent was obtained from each of the subjects before their participation in the study.

The Mini Mental State Examination (MMSE) is widely used as a global test of cognitive function to screen for dementia and assesses orientation, immediate memory, attention and calculation, short-term recall, and language [9]. Based on the original MMSE, we developed a modified and culturally adapted MMSE according to Indonesian subjects. The MMSE (score range, 0-30) is used as a screening test for cognitive impairment. The Recall test (score range, 0-3) is a sub-item of the IMMSE that evaluates the impairment of recent memory. Subjects were instructed to recall three unrelated objects that they were previously instructed to remember. The MMSE was carried out by two interviewers who were formally trained by a psychiatrist. Cognitive impairment was classified as an MMSE score of less than 24 [9, 10].

A dental examination was carried out by two dentists calibrated regarding the techniques, with the dentist and the subject in a sitting position under artificial lighting. The number of teeth was recorded for each subject. The remaining teeth were defined as healthy, carious, or treated teeth, inclusive of completely erupted third molars. Severe carious teeth and very loose teeth that were indicated for extraction were excluded as remaining teeth. We categorized the remaining teeth as twenty teeth or more, eleven to nineteen, ten teeth or less.

We classified that having the status of hypertension when a participant's having systolic blood pressure of 140 mm Hg or greater and/or diastolic blood pressure of 90 mm Hg or greater. We categorized participant's non-fasting glucose level as 200 mg/dL or over and below than 200 mg/dL.

Statistical analyses were performed using Confidence Interval Analysis Software for Windows (ver. 2.0, Trevor Bryant, University of Southampton). Logistic regression analysis was carried out with low MMSE score as dependent variables and other variables were used as independent variables. The number of remaining teeth was used as continuous variables in one model and categorical variables in the other model.

## RESULTS AND DISCUSSION

### Results

All of the 208 participants aged 60 years old and above (36 men and 172 women) have completed all the examination processes. The subjects were divided into two groups based on their MMSE scores. The demographic characteristics of subjects, the socioeconomic characteristics and the number of remaining teeth are shown in Table 1.

The mean age (SD) of male and female subjects were 74.4 (8.5) and 66.8 (6.9) years old, respectively. The mean MMSE score (SD) of subjects with low MMSE score ( $\leq 23$ ) and high MMSE score ( $\geq 24$ ) were 20.1 (8.8) and 27.0 (2.0), respectively. The proportions of men and women having low MMSE were found 25% (9/36) and 25.6% (44/172), respectively. Subjects living in the rural area were more likely to have a lower MMSE score. The proportions of urban and rural elders having low MMSE were found 20.7% (32/156) and 39.6% (21/53) respectively. Most of subjects with low MMSE were having 6 years or lower education experience (37.8%). The proportion of subjects with low MMSE among 7 - 12 years and  $\geq 13$  years educational experience groups were 19.4% (21/108) and 5.6% (1/18), respectively. Monthly allowances among subjects with low MMSE was found as low as IDR 512,903 while in the high MMSE group was found as high as IDR 922,288.

The mean (SD) number of remaining teeth in subjects with low MMSE and high MMSE were 14.65 (7.96) and 21.5 (8.02), respectively. In the logistic regression analysis, odds ratio (OR) of cognitive impairment (MMSE score  $\leq 23$ ) were 6.34 (95% CI, 2.76 to 14.55) and 2.82 (95% CI, 1.30 to 6.10) for subjects who had remaining  $\leq 10$  teeth and those who had remaining 11 - 19 teeth, respectively, when compared with subjects who had remaining  $\geq 20$  teeth.

### Discussion

The result of this community-based survey revealed that the

**Table 1:** Distribution of demographic, socioeconomic, and number of remaining teeth status.

Parameter	Low MMSE ( $\leq 23$ )		High MMSE ( $\geq 24$ )	
	Mean (SD)	n (%)	Mean (SD)	n (%)
<b>Age (years), mean (SD)</b>	73.45	(8.92)	66.29	(5.38)
<b>Sex, number (%)</b>				
males	9	(25.0)	27	(75.0)
females	44	(25.6)	128	(74.4)
<b>Region, number (%)</b>				
urban	32	(15.38)	123	(59.13)
rural	21	(10.10)	32	(15.38)
<b>Length of education, number (%)</b>				
$\geq 13$ yrs	1	(5.6)	17	(94.4)
7 - 12 yrs	21	(19.4)	87	(80.6)
$\leq 6$ yrs	31	(37.8)	51	(62.2)
<b>Allowances (IDR/month), mean (SD)</b>	512,903	(507,452)	1,541,389	(922,288)
<b>Number of remaining teeth, mean (SD)</b>	14.65	(7.96)	21.50	(8.02)

**Abbreviations:** IDR: Indonesian Rupiah; MMSE: Mini Mental State Examination; SD: Standard Deviation

prevalence of a low MMSE score was significantly associated with low number of remaining teeth. These results are consistent with those of previous studies demonstrating that tooth loss was associated with decreased cognitive function [5,7,8]. Many factors may contribute to cognitive impairment in elders, including cardiovascular disease, education and social economy status [10].

The other main findings of this study are that a low level of education and a low income were significantly associated with low MMSE score and increased risk of dementia. Our results are in agreement with previous findings in a study of education and dementia in a Japanese population [7]. Different mechanisms have been suggested to explain this association. Katzman [11] proposed that education might increase brain reserves by increasing synaptic density in the neocortical association cortex. Stern et al. extended the cognitive reserve hypothesis and took into account the possible beneficial influence of mental activity throughout the entire life span, where occupational attainment along with level of education could influence the risk of dementia's disease [12]. It is known that education is an indicator of socioeconomic status. This socioeconomic status acting over the life course have been found to affect health and risk of cardiovascular disease. Cumulative economic hardship has been found to lead to poorer cognitive functioning and dementia.

The association between loss of molars and spatial memory deficits was first described in an animal study using senile rats [13]. In that study, after molar extraction, rats were maintained with diet powder and were subjected to behavioral testing to evaluate their spatial memory. Animals with extracted molar teeth showed worse behavioral performance compared to control animals maintained with solid diet. Mastication is the main function of stomatognathic system and the relationship between mastication and the central nervous system (CNS), especially the cognitive area, is still the subject of several studies. Numerous studies have revealed an association between mastication and cognitive functions [10,13-16]. The possible associations between masticatory hypofunction and cognitive insufficiency have been demonstrated in several animal studies [16-19]. Rats in those studies were induced masticatory hypo function by performing extraction or occlusal reduction of the molar teeth rather than a long-term soft diet [16,17,19]. The occlusal hypo function caused degenerative changes in their periodontal mechanoreceptors [20] suggesting a suppression of sensory stimulation from the periodontal ligament during chewing. In those experiments, animals showed poor performance on memory and learning evaluation.

Histologically, the hippocampus, the main CNS region, that affected by masticatory hypofunction was reported by several studies [16,21]. The hippocampus is included in some kinds of memory, especially which relates to details and precise events [16, 21]. Recently, an animal study using senescence-accelerated mouse prone 8 (SAMP8) examined whether reduced mastication from a young age mice affects hippocampal-dependent cognitive function [22] (Table 2). This study found that loss of masticatory function in early life causes malnutrition and chronic stress and impairs the ability to recognize novel objects. Hyper activation and lateralized rotational behavior are commonly observed with dysfunction of the dopaminergic system, therefore,

**Table 2:** Logistic regression model for predicting lower MMSE scores ( $\leq 23$ ).

Variables	OR	95% CI	P value
<b>Age</b>			
$\leq 65$ years (reference)	1.00		
$\geq 66$ years	8.86	3.76 to 20.87	< 0.05
<b>Sex</b>			
males (reference)	1.00		
females	1.03	0.45 to 2.36	> 0.05
<b>Region</b>			
Urban (reference)	1.00		
Rural	2.52	1.29 to 4.95	< 0.05
<b>Length of education</b>			
13 years or over (reference)	1.00		
7 - 12 years	4.10	0.52 to 32.60	> 0.05
6 years or under	10.33	1.31 to 81.53	< 0.05
<b>Allowances (IDR/month)</b>			
$\geq 1,500,000$ (reference)	1.00		
< 1,500,000	3.01	1.56 to 5.82	< 0.05
<b>Blood pressure</b>			
Normotension (reference)	1.00		
Hypertension	1.38	0.68 to 2.81	< 0.05
<b>Non fasting glucose level (mg/dL)</b>			
< 200 (reference)	1.00		
$\geq 200$	0.17	0.02 to 1.29	> 0.05
<b>Number of remaining teeth</b>			
$\geq 20$ (reference)	1.00		
11-19	2.82	1.30 to 6.10	< 0.05
$\leq 10$	6.34	2.76 to 14.55	< 0.05

**Abbreviations:** CI: Confidence Interval; IDR: Indonesian Rupiah; Mg/dL: Milligram per Deciliter; OR: Odds Ratio.

reduced masticatory function may deplete the mesolimbic and mesocorticolimbic dopaminergic systems to impair the cognitive functions of selective attention and recognition memory in the prefrontal cortex and the hippocampus [22-24].

Masticatory dysfunction also leads to various morphologic changes in the hippocampus and cerebral cortex, e.g., the molar less condition and soft-diet feeding decrease the brain cells including synapses in the hippocampus and parietal cortex [25], as well as neurotrophic receptor expression in hippocampal CA1 and CA3 regions [26]. Hypertrophy and hyperplasia of glial fibrillary acid protein-labeled astrocytes in the CA1 region [27, 28] and suppressed cell proliferation in the hippocampal dentate gyrus [29,30] also occur in mice with learning deficits. These behavioral and hippocampal morphologic changes are similar to the changes associated with advanced age [31,32] and masticatory dysfunction may therefore accelerate the aging process in the hippocampus.

Neuroimaging clinical trials using functional MRI and positron emission tomography indicate that several regions of the brain

are activated during mastication, including the cortex, thalamus, striatum, and cerebellum [33-35]. Functional MRI evaluation on neuronal activity in the brain indicated that chewing activity increases the blood oxygen level-dependent signals in the right premotor cortex, precuneus, thalamus, hippocampus and inferior parietal lobe [36]. Furthermore, chewing activity was reported significantly enhances memory in aged subjects [37]. Cerebellar functions also seem to be influenced by masticatory activity [34]. Recently, Quintero and others [38] have reported that the functional connections between cerebellum and cortical areas during masticatory activity [36-38]. It has been well documented that the cerebellum is responsible for motor planning, including the force needed to perform motor tasks. However, it is not clear yet what is the mechanism of reduced input activity due to masticatory dysfunction differ from the effects of a reduction in other types of sensory stimuli in the aging hippocampus. To elucidate the mechanisms underlying the specific effects of masticatory dysfunction in aging hippocampus, it will be necessary to compare the impact of mastication and environmental enrichment or physical exercise on hippocampal function.

## CONCLUSION

In summary, there are significant relationships were found between the number of remaining teeth and cognitive function. Edentulism may be predictors of dementia late in life.

## ACKNOWLEDGEMENTS

This work was supported by Universitas Trisakti, Jakarta Indonesia. The authors declare no potential conflicts of interest with respect to the authorship and/or publication of this article. We thank Dr. Widijanto Sudhana, M. Kes, Faculty of Dentistry, Universitas Trisakti for constructive statistical comments which have assisted us in preparing this article.

## REFERENCES

1. WHO. Dementia: a public health priority. Geneva: World Health Organization; 2012.
2. Ferri CP, Prince M, Brayne C, Brodaty H, Fratiglioni L, Ganguli M, et al. Global prevalence of dementia: a Delphi consensus study. *Lancet*. 2005; 366: 2112-2117.
3. International Alzheimer's Disease. The Prevalence of Dementia Worldwide. 2008.
4. Stein PS, Desrosiers M, Donegan SJ, Yepes JF, Kryscio RJ. Tooth loss, dementia and neuropathology in the Nun study. *J Am Dent Assoc*. 2007; 138: 1314-1322.
5. Okamoto N, Morikawa M, Okamoto K, Habu N, Iwamoto J, Tomioka K, et al. Relationship of tooth loss to mild memory impairment and cognitive impairment: findings from the Fujiwara-kyo study. *Behav Brain Funct*. 2010; 6: 77.
6. Kaye EK, Valencia A, Baba N, Spiro A 3rd, Dietrich T, Garcia RI. Tooth loss and periodontal disease predict poor cognitive function in older men. *J Am Geriatr Soc*. 2010; 58: 713-718.
7. Saito Y, Sugawara N, Yasui-Furukori N, Takahashi I, Nakaji S, Kimura H. Cognitive function and number of teeth in a community-dwelling population in Japan. *Ann Gen Psychiatry*. 2013; 12: 20.
8. Park H, Suk SH, Cheong JS, Lee HS, Chang H, Do SY, Kang JS. Tooth loss may predict poor cognitive function in community-dwelling adults without dementia or stroke: the PRESENT project. *J Korean Med Sci*. 2013; 28: 1518-1521.
9. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975; 12: 189-198.
10. Qiu C, De Ronchi D, Fratiglioni L. The epidemiology of the dementias: an update. *Curr Opin Psychiatry*. 2007; 20: 380-385.
11. Katzman R. Education and the prevalence of dementia and Alzheimer's disease. *Neurology*. 1993; 43: 13-20.
12. Stern Y, Gurland B, Tatemichi TK, Tang MX, Wilder D, Mayeux R. Influence of education and occupation on the incidence of Alzheimer's disease. *JAMA*. 1994; 271: 1004-1010.
13. Kato T, Usami T, Noda Y, Hasegawa M, Ueda M, Nabeshima T. The effect of the loss of molar teeth on spatial memory and acetylcholine release from the parietal cortex in aged rats. *Behav Brain Res*. 1997; 83: 239-242.
14. Miura H, Miura K, Mizugai H, Arai Y, Umenai T, Isogai E. Chewing ability and quality of life among the elderly residing in a rural community in Japan. *J Oral Rehabil*. 2000; 27: 731-734.
15. Miura H, Yamasaki K, Kariyasu M, Miura K, Sumi Y. Relationship between cognitive function and mastication in elderly females. *J Oral Rehabil*. 2003; 30: 808-811.
16. Scherder E, Posthuma W, Bakker T, Vuijk PJ, Lobbezoo F. Functional status of masticatory system, executive function and episodic memory in older persons. *J Oral Rehabil*. 2008; 35: 324-336.
17. Ono Y, Yamamoto T, Kubo KY, Onozuka M. Occlusion and brain function: mastication as a prevention of cognitive dysfunction. *J Oral Rehabil*. 2010; 37: 624-640.
18. Kato T, Usami T, Noda Y, Hasegawa M, Ueda M, Nabeshima T. The effect of the loss of molar teeth on spatial memory and acetylcholine release from the parietal cortex in aged rats. *Behav Brain Res*. 1997; 83: 239-242.
19. Kushida S, Kimoto K, Hori N, Toyoda M, Karasawa N, Yamamoto T. Soft-diet feeding decreases dopamine release and impairs aversion learning in Alzheimer model rats. *Neurosci Lett*. 2008; 439: 208-211.
20. Ekuni D, Tomofuji T, Irie K, Azuma T, Endo Y, Kasuyama K, et al. Occlusal disharmony increases amyloid- $\beta$  in the rat hippocampus. *Neuromolecular Med*. 2011; 13: 197-203.
21. Yamamoto T, Hirayama A. Effects of soft-diet feeding on synaptic density in the hippocampus and parietal cortex of senescence-accelerated mice. *Brain Res*. 2001; 902: 255-263.
22. Muramoto T, Takano Y, Soma K. Time-related changes in periodontal mechanoreceptors in rat molars after the loss of occlusal stimuli. *Arch Histol Cytol*. 2000; 63: 369-380.
23. Squire LR, Stark CE, Clark RE. The medial temporal lobe. *Annu Rev Neurosci*. 2004; 27: 279-306.
24. Kawahata M, Ono Y, Ohno A, Kawamoto S, Kimoto K, Onozuka M. Loss of molars early in life develops behavioral lateralization and impairs hippocampus-dependent recognition memory. *BMC Neurosci*. 2014; 15: 4.
25. Yamamoto T, Hirayama A. Effects of soft-diet feeding on synaptic density in the hippocampus and parietal cortex of senescence-accelerated mice. *Brain Res*. 2001; 902: 255-263.
26. Yamazaki K, Wakabayashi N, Kobayashi T, Suzuki T. Effect of tooth loss on spatial memory and trkB-mRNA levels in rats. *Hippocampus*. 2008; 18: 542-547.
27. Onozuka M, Watanabe K, Nagasaki S, Jiang Y, Ozono S, Nishiyama

- K, et al. Impairment of spatial memory and changes in astroglial responsiveness following loss of molar teeth in aged SAMP8 mice. *Behav Brain Res.* 2000; 108: 145-155.
28. Watanabe K, Tonosaki K, Kawase T, Karasawa N, Nagatsu I, Fujita M, et al. Evidence for involvement of dysfunctional teeth in the senile process in the hippocampus of SAMP8 mice. *Exp Gerontol.* 2001; 36: 283-295.
29. Aoki H, Kimoto K, Hori N, Toyoda M. Cell proliferation in the dentate gyrus of rat hippocampus is inhibited by soft diet feeding. *Gerontology.* 2005; 51: 369-374.
30. Kubo KY, Sasaguri K, Ono Y, Yamamoto T, Takahashi T, Watanabe K, et al. Chewing under restraint stress inhibits the stress-induced suppression of cell birth in the dentate gyrus of aged SAMP8 mice. *Neurosci Lett.* 2009; 466: 109-113.
31. Kitraki E, Bozas E, Philippidis H, Stylianopoulou F. Aging-related changes in IGF-II and c-fos gene expression in the rat brain. *Int J Dev Neurosci.* 1993; 11: 1-9.
32. DeToledo-Morrell L GYaMF. Age-dependent alteration in hippocampal synaptic plasticity: relation to memory disorders. *Neurobiol Aging.* 1993; 14: 441-446.
33. Momose T, Nishikawa J, Watanabe T, Sasaki Y, Senda M, Kubota K, et al. Effect of mastication on regional cerebral blood flow in humans examined by positron-emission tomography with  $\mu$ O-labelled water and magnetic resonance imaging. *Arch Oral Biol.* 1997; 42: 57-61.
34. Onozuka M, Fujita M, Watanabe K, Hirano Y, Niwa M, Nishiyama K, et al. Mapping brain region activity during chewing: a functional magnetic resonance imaging study. *J Dent Res.* 2002; 81: 743-746.
35. Onozuka M, Fujita M, Watanabe K, Hirano Y, Niwa M, Nishiyama K, et al. Age-related changes in brain regional activity during chewing: a functional magnetic resonance imaging study. *J Dent Res.* 2003; 82: 657-660.
36. Hirano Y, Obata T, Kashikura K, Nonaka H, Tachibana A, Ikehira H, et al. Effects of chewing in working memory processing. *Neurosci Lett.* 2008; 436: 189-192.
37. Onozuka M, Tachibana A, Kim W, Ono Y, Sasaguri K, Kubo K, Niwa M, et al. Interactions between chewing and brain activities in human. *Novel Trends in Brain Science.* 2007; 99-113.
38. Quintero A, Ichesco E, Schutt R, Myers C, Peltier S, Gerstner GE. Functional connectivity of human chewing: an fMRI study. *J Dent Res.* 2013; 92: 272-278.

**Cite this article**

Asia A, Kusdhany L, Rahardjo A, Bachtiar A (2015) Association between Tooth Loss and the Development of Dementia on the Indonesian Elders. *JSM Dent* 3(1): 1050.

# Association between Tooth Loss and the Development of Dementia on the Indonesian Elders

---

## ORIGINALITY REPORT

---

**22%**

SIMILARITY INDEX

**12%**

INTERNET SOURCES

**19%**

PUBLICATIONS

**10%**

STUDENT PAPERS

---

## MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

3%

★ [www.pubfacts.com](http://www.pubfacts.com)

Internet Source

---

Exclude quotes  On

Exclude matches  < 15 words

Exclude bibliography  On

# Association between Tooth Loss and the Development of Dementia on the Indonesian Elders

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---