

SERTIFIKAT

Kementerian Riset dan Teknologi/
Badan Riset dan Inovasi Nasional



Petikan dari Keputusan Menteri Riset dan Teknologi/
Kepala Badan Riset dan Inovasi Nasional
Nomor 200/M/KPT/2020
Peringkat Akreditasi Jurnal Ilmiah Periode III Tahun 2020
Nama Jurnal Ilmiah

Jurnal Ilmiah Pendidikan Fisika Al-Biruni

E-ISSN: 2503023X

Penerbit: Program Studi Pendidikan Fisika, UIN Raden Intan Lampung

Ditetapkan sebagai Jurnal Ilmiah

TERAKREDITASI PERINGKAT 2

Akreditasi Berlaku selama 5 (lima) Tahun, yaitu
Volume 9 Nomor 1 Tahun 2020 sampai Volume 13 Nomor 2 Tahun 2024

Jakarta, 23 December 2020

Menteri Riset dan Teknologi/
Kepala Badan Riset dan Inovasi Nasional
Republik Indonesia,



Bambang P. S. Brodjonegoro
Bambang P. S. Brodjonegoro



COLLABORATE WITH



USER

Username

Password

Remember me

Login

JOURNAL CONTENT

Search

Search Scope
All

Search

Browse

Home > Vol 11, No 2 (2022)

Jurnal Ilmiah Pendidikan Fisika Al-Biruni

Jurnal ilmiah pendidikan fisika Al-Biruni [JIPF Al-Biruni] is a multidisciplinary and a medium of communication for researchers, academicians, and practitioners that provides a means for sustained discussion of relevant issues that fall within the focus and scopes of the journal which examined empirically.

JIPF Al-Biruni covers issues such as the **improvement of physics learning techniques** including the development of instruments of physics evaluation, the development of physics instructional media, and the development of learning model of physics by considering local wisdom and being enriched with the principle of religion and culture. **Theoretical physics and current issue of applied physics** are also welcome. However, the article related and integrated to the value of religion and culture will be prioritized. **Development and educational psychology in context of learning physics** such as: learning physics in early education, learning physics in primary and secondary education, and learning physics in higher education

The print version of JIPF Al-Biruni was first published in 2012 then followed by the online version in 2014.



Journal title	Jurnal ilmiah pendidikan fisika Al-Biruni
Initials	JIPF Al-Biruni
Accreditation	Sinta Rank 2 by Ministry of Research, Technology, and Higher Education of the Republic of Indonesia
Frequency	2 issues per year (April and October)
DOI	Prefix 10.24042 by Crossref
Online ISSN	2503-023X
Editor-in-chief	Antomi Saregar
Publisher	Physics Education Department, Universitas Islam Negeri Raden Intan Lampung in collaboration with Perkumpulan Pendidik IPA Indonesia (PPII) and Physics Society of Indonesia (PSI)
Citation Analysis	Google Scholar SINTA DIMENSION
Submissions	This journal welcomes submissions all year round. Submissions are reviewed upon receipt. The template manuscript can be downloaded Here . Manuscript Published is written in standard English since Volume 7 Number 1, 2018.

- People
 - Editorial Team
 - Reviewer
 - Contact
- Policies
 - Journal History
 - Focus and Scope
 - Peer Review Process
 - Publication Frequency
 - Open Access Policy
 - Digital Preservation
 - Deposit Policy
 - Publication Ethics and Misconduct
 - Screening for Plagiarism
 - Withdrawal of Manuscript
 - Correction and Retraction
 - Indexing and Abstracting
 - Article Processing Charge
 - Revenue Sources
- Submissions
 - Author Guidelines
 - Copyright Notice

Tim Uji Kesamaan



COLLABORATE WITH



USER

Username

Password

Remember me

JOURNAL CONTENT

Search

Search Scope

Browse

- ▶ By Issue
- ▶ By Author
- ▶ By Title
- ▶ Other Journals

INFORMATION

- ▶ For Readers
- ▶ For Authors

Home > About the Journal > Editorial Team

Editorial Team

Editor in Chief

Antomi Saregar, (Scopus ID: 57201321267, Subject area: Physics Education) Universitas Islam Negeri Raden Intan Lampung, Indonesia

Advisory Editorial Board

Konstantinos Ravanis, (Scopus ID: 8710602700, Subject area: Physics Education, Science Education) University of Patras, Greece

Jean Marie Boilevin, (Scopus ID: 24470928200, Subject area: Physics Education), Centre de Recherche sur l'Education, les Apprentissages et la Didactique - CREAD, Université de Bretagne Occidentale., France

Adel Koth, (Scopus ID: 55614316800, Subject area: Physics, Hydro-geophysics, Near Surface Geophysics, Applied Geophysics) Helwan University, Cairo, Egypt

Mitra Djamal, (Scopus ID: 6506355194, Subject area: Applied Physics) Institut Teknologi Sumatera and Chief of Physics Society of Indonesia, Indonesia

Andrew Heckler, (Scopus ID: 6602388258, Subject area: Physics Education) Ohio State University, Department of Physics, Columbus, United States

Warsito Warsito, (Scopus ID: 37079631800, Subject area: Physics and Astronomy; Materials Science; Earth and Planetary Sciences) Universitas Lampung, Indonesia

Cari Cari, (Scopus ID: 33267518100, Subject area: Physics Education) Universitas Negeri Sebelas Maret, Indonesia, Indonesia

Anto Sulaksono, (Scopus ID: 6507369203 Subject area: Physics and Astronomy) Universitas Indonesia, Indonesia

Editorial Board

Rofiqul Umam, (Scopus ID: 57201308953) Faculty of Science and Technology, Kwansai Gakuin University, Japan

Vishal R Panse, Ph. D., Late.B.S.Arts, Prof.N.G.Science & A.G.Commerce College, Sakarkherda, India

Ishwar Shrawan Mohurley, (Solid State Physics and Relativity) Shri Dnyanesh Mahavidyalaya Nawargaon, India

Abdurrahman Abdurrahman, (Scopus ID: 57006623600, Subject area: Physics Education) Universitas Lampung, Indonesia

Hartono Bancong, (Scopus ID: 57200101660, Subject area: Physics Education), Universitas Muhammadiyah Makassar, Indonesia

Ali Rahmat, (Scopus ID: 57189066685; Applied Physics for renewable energy) Lembaga Ilmu Pengetahuan Indonesia [The Indonesian Institute of Sciences], Indonesia

Happy Komikesari, (Scopus ID: 57208305928, Subject area: Physics Education) Universitas Islam Negeri Raden Intan Lampung, Indonesia

Achmad Samsudin, (Scopus ID: 57191537500, Subject area: Physics Education) Universitas Pendidikan Indonesia, Indonesia

Dwi Yulianti, (Scopus ID: 57194041989) Universitas Negeri Semarang, Indonesia


Rahma Diani, (Scopus ID: 57205348300) Universitas Islam Negeri Raden Intan Lampung, Indonesia

Administrator

Nur Endah Susilowati, (Scopus ID: 57208302422, Subject area: Physics Education) Universitas Islam Negeri Raden Intan Lampung, Indonesia

IT Support

Adyt Anugrah, (Scopus ID: 57211393785; Subject area: Physics Education) Universitas Islam Negeri Raden Intan Lampung, Indonesia



People
Editorial Team
Reviewer
Contact

Policies
Journal History
Focus and Scope
Peer Review Process
Publication Frequency
Open Access Policy
Digital Preservation
Deposit Policy
Publication Ethics and Misconduct
Screening for Plagiarism
Withdrawal of Manuscript
Correction and Retraction
Indexing and Abstracting
Article Processing Charge
Revenue Sources

Submissions
Author Guidelines
Copyright Notice
Privacy Statement

Information
For Readers
For Authors
For Librarians





Article template

COLLABORATE WITH



USER

Username

Password

Remember me

JOURNAL CONTENT

Search

Search Scope
 All

Browse

- ▶ By Issue
- ▶ By Author
- ▶ By Title
- ▶ Other Journals

INFORMATION

- ▶ For Readers

Home > About the Journal > People

People

Reviewers

- Agus Purwanto, (Scopus ID: 56547539300, Subject area: Physics and Astronomy; Mathematics) ITS Surabaya, Indonesia
- Lilia Halim, (Scopus ID: 14624924300, Subject area: Physics Education) Universiti Kebangsaan Malaysia, Malaysia
- Fatin Aliah Phang, (Scopus ID: 57192376607, Subject area: Physics, Physics and Astronomy, Computer Science) Universiti Teknologi Malaysia
- Muhammad Mus'ab Anas, (Scopus ID: 56945771600, Subject area: Physics Education, Quantum Physics) Universiti Sains Islam Malaysia
- George Sahaya Dennish Babu, (Scopus ID: 57193324733), Chettinad College of Engineering and Technology, India
- Irzaman Irzaman, (Scopus ID: 7409571162; Subject area: Materials Physics [BioSensor, Thin Film, Bioenergy]) Institut Pertanian Bogor
- Zuhdan Kun Prasetyo, (Scopus ID: 57190379913, Subject area: Physics Education) Universitas Negeri Yogyakarta, Indonesia
- Agus Setiawan, (Scopus ID: 6701505999, Subject area: Physics, Materials Science) UPI Bandung, Indonesia
- Andi Suhandi, (Scopus ID: 35103628800, Subject area: Physics, Physics Education) Universitas Pendidikan Indonesia, Indonesia
- Mehmet Fatih Tasar, (Scopus ID: 13410804800) Gazi University, Turkey
- Putut Marwoto, (Scopus ID: 36904245100, Subject area: Physics and Astronomy) Universitas Negeri Semarang, Indonesia
- Ketut Suma, (Scopus ID: 57202671029, Subject area: Physics Education) Universitas Pendidikan Ganesha, Indonesia
- Abdurrouf Abdurrouf, (Scopus ID: 12646267900, Subject area: Physics and Material) Universitas Brawijaya, Indonesia
- Soeharto Soeharto, (Scopus ID: 57210115604) University of Szeged, Hungary
- Doni Andra, (Scopus ID: 6507316610, Subject area: Physics, Physics Education) University of Lampung, Indonesia, Indonesia
- Dadi Rusdiana, (Scopus ID: 56070183500, Subject area: Physics Education) Universitas Pendidikan Indonesia, Indonesia
- Dadan Rosana, (Scopus ID: 57195052389, Subject area: Physics Education; Physics) Universitas Negeri Yogyakarta, Indonesia
- Agus Suyatna, (Scopus ID: 557199237734, Subject area: Science Education, Physics Education) Universitas Lampung, Indonesia
- Muqoyyanah Muqoyyanah, (Scopus ID: 57200148823, Subject area: Nanomaterial Physics) Badan Riset dan Inovasi Nasional, Indonesia
- John Rafafy Batlolona, (Scopus ID: 57192589179, Subject area: Physics Education) Universitas Pattimura, Indonesia



People
Editorial Team
Reviewer
Contact

Policies
Journal History
Focus and Scope
Peer Review Process
Publication Frequency
Open Access Policy
Digital Preservation
Deposit Policy
Publication Ethics and Misconduct
Screening for Plagiarism
Withdrawal of Manuscript
Correction and Retraction
Indexing and Abstracting
Article Processing Charge
Revenue Sources

Submissions
Author Guidelines
Copyright Notice
Privacy Statement

Information
For Readers
For Authors
For Librarians



<p>Specific Open-Ended Assessment: Assessing Students' Critical Thinking Skill on Kinetic Theory of Gases</p> <p> <i>Riki Perdana, Riwayani Riwayani, Jumadi Jumadi, Dadan Rosana, Soeharto Soeharto</i></p> <p> 10.24042/jipfalbiruni.v8i2.3952  Abstract views : 1014</p>	127-140 
<p>Determination of Groundwater Using Geoelectric Methods: Schlumberger Configuration in Rokan Hulu Regency</p> <p> <i>Yeza Febriani, Riza Ari Rohman, Azmi Asra, Mohammad Apriniyadi, Donata Nariswari Wahyu Wardani</i></p> <p> 10.24042/jipfalbiruni.v8i2.3918  Abstract views : 1151</p>	141-152 
<p>How is Students' Creative Thinking Skills? An Ethnoscience Learning Implementation</p> <p> <i>Ahmad Khoiri, Nulingafan Nulingafan, Widha Sunarno, Sajidan Sajidan</i></p> <p> 10.24042/jipfalbiruni.v8i2.4559  Abstract views : 1838</p>	153-163 
<p>Academic Achievement and Conceptual Understanding of Electrodynamics: Applications Geoelectric Using Cooperative Learning Model</p> <p> <i>Rizqi Prastowo, Syamsul Huda, Rafiqul Umam, Kittisak Jemsittiparsert, Andika Eko Prasetyo, Hasan Said Tortop, Muhamad Syazali</i></p> <p> 10.24042/jipfalbiruni.v8i2.4614  Abstract views : 548</p>	165-175 
<p>Murai Batu's (<i>Copsychus malabaricus</i>) Peak Frequency Sound: The Impact toward Stomatal Pores of The Cayenne Pepper (<i>Capsicum frutescens L</i>) Leaves</p> <p> <i>Eka Sobiatin, Herianto Herianto, Nur Khosiyatun, Mirra Fatharani, Heru Kuswanto</i></p> <p> 10.24042/jipfalbiruni.v8i2.4137  Abstract views : 661</p>	177-185 
<p>Project-Based (PjBL) and Guided Inquiry Learning: Students' Response to Rectilinear and Circular Motion Phenomena</p> <p> <i>Rismawati Rismawati, Sarwanto Sarwanto, Budiyo Saputra</i></p> <p> 10.24042/jipfalbiruni.v8i2.4422  Abstract views : 710</p>	187-196 
<p>Local wisdom-Based Electronic Book on Newton's Law</p> <p> <i>Trisya Afidah Sukma, Mundilarto Mundilarto, Nadia Darma Putri</i></p> <p> 10.24042/jipfalbiruni.v8i2.4368  Abstract views : 556</p>	197-209 
<p>Digital Story Conceptual Change Oriented (DSCC) to Reduce Students' Misconceptions in Physics</p> <p> <i>Yudi Kurniawan, Riski Mulyani, Shahala Nassim</i></p> <p> 10.24042/jipfalbiruni.v8i2.4596  Abstract views : 706</p>	211-220 
<p>Students' Creative Thinking Skills on Heat Phenomena Using POGIL Learning Model</p> <p> <i>Rinta Dian Pratiwi, Ashadi Ashadi, Sukarmin Sukarmin, Dewanto Harjunowibowo</i></p> <p> 10.24042/jipfalbiruni.v8i2.4629  Abstract views : 651</p>	221-231 
<p>Android-Based Physics Learning Media Integrated Landslide Disaster</p> <p> <i>Ulfaturrone Nur Labibah, Insih Wilujeng, Syazana Sulaiman, Lia Rahmawati</i></p> <p> 10.24042/jipfalbiruni.v8i2.4695  Abstract views : 899</p>	233-240 
<p>Mobile Phone Utilization for Learning: Physics Teachers' Perception</p> <p> <i>Agus Suyatna</i></p> <p> 10.24042/jipfalbiruni.v8i2.4840  Abstract views : 594</p>	241-248 
<p>Inquiry-Scaffolding Learning Model: Its Effect on Critical Thinking Skills and Conceptual Understanding</p> <p> <i>Wartono Wartono, Yohana Fransiska Alfroni, John Rafafy Batlolona, Natcha Mahapoonyanont</i></p> <p> 10.24042/jipfalbiruni.v8i2.4214  Abstract views : 702</p>	249-259 



Determination of Groundwater Using Geoelectric Methods: Schlumberger Configuration in Rokan Hulu Regency

Yeza Febriani^{*1}, Riza Ari Rohman², Azmi Azra³, Mohammad Apriniyadi⁴, Donata Nariswari Wahyu Wardani⁵

^{1,2,3}Physics Study Programme, Faculty of teacher training and education, Pasir Pengaraian University, Rokan Hulu Regency 28557, Riau Province, Indonesia;

⁴Geology Technic Programme, Universitas Trisakti, Jakarta Barat, Jakarta, Indonesia;

⁵Groundwater management Programme, Faculty of Geoscience, Geoengineering and Mining, Freiberg University of Mining and Technology, Akademiestraße 609599 Freiberg, Germany.

*Corresponding Address: yezafebriani@upp.ac.id

Article Info

Article history:

Received: April 1st, 2019

Accepted: October 20th, 2019

Published: October 30th, 2019

Keywords:

aquifer;
 geoelectric;
 groundwater;
 resistivity;
 schlumberger.

ABSTRACT

This study aimed at detecting the presence of groundwater in Masda Makmur Village, Rambah Samo District, Rokan Hulu Regency, using the Schlumberger configuration geoelectric method. The study consisted of 3 sounding points within every 100 m sounding track length. The measurement data is processed using IP2WIN software and Progress to see layer data below the ground level based on the value of the resistivity type. The results showed that the estimated groundwater could be found on track one with a kind of resistivity value of 7,44 Ωm at a depth of 15-22 m. Alleged groundwater can be found on the second track with a type of resistivity value of 75,73 Ωm at a depth of 13-18 m, and the estimated groundwater can be located on the third track with a kind of resistivity value of 82,52 Ωm , 93,26 Ωm . The deeper and the lower resistivity value shows that the layer has the potential as a carrier layer of groundwater (aquifer). Sounding 1 and 2 have the potential to make bore wells which are thought to be depressed aquifer. The results of this study can be used further for mapping the location of community boreholes

© 2019 Physics Education Department, UIN Raden Intan Lampung, Indonesia.

INTRODUCTION

Water is essential for life on earth (Saranga et al., 2016). Water consists of two atoms of hydrogen and one atom of oxygen (H_2O). Living things cannot live without water. The increasing population requires a sufficient amount of water. In Indonesia, especially Rokan Hulu Regency, population growth is increasing every year (Bappeda Rokan Hulu, 2015).

In 2016, the percentage of population growth rate in the Rokan Hulu Regency was 0.27%, 2017 experienced an increase of 0.09%, and in 2018 there was an increase of

0.05%. For Rambah Samo District, Rokan Hulu Regency in 2016 the population growth rate was 1,36%, 2017 increased by 1,82% and in 2018 again experienced a significant increase of 2,05% (BPS Rokan Hulu, 2017). The higher of the population growth rate, the more need for clean water by the community. Groundwater is one source of life necessities for living things on earth (Hanifa, Sota, & Siregar, 2016; Seha & Aziz, 2016). Groundwater is all water found in layers containing water (aquifers) below the surface of the soil. The role of groundwater is increasingly essential to

How to cite

Febriani, Y., Rohman, R. A., Asra, A., Apriniyadi, M., & Wardani, D. N. W. (2019). Determination of Groundwater Using Geoelectric Methods: Schlumberger Configuration in Rokan Hulu Regency. *Jurnal Ilmiah Pendidikan Fisika Al-Biruni*, 8(2), 141-152.

become the primary water source to fulfill the basic needs of many people (Jusuf, 2015; Sedana, As'ari, & Tanauma, 2015).

Specific methods can detect the presence of groundwater in the subsurface layer (Devi, Israil, Anbalagan, & Gupta, 2017; Heradian & Arman, 2015; Wijaya, 2015). One of which is by using the geoelectric resistivity method. This method is the most popular method of investigating groundwater from the surface of the earth (Bakri, 2019; Bashir, Izham, & Main, 2014; El-Hameed, El-Shayeb, El-Araby, & Hegab, 2017; Helaly, 2017; Hewaidy et al., 2015; Jusuf, 2015; Mohamaden, 2016). The geoelectric method is a method that studies the nature of electricity on the planet by detecting it on the earth's surface (Hakim & Manrulu, 2017).

Several studies on the detection of groundwater estimation using Schlumberger configuration have been carried out, including editions (Edisar, 2013) of data obtained, namely zoning of free aquifers and depressed aquifers. Sehad and Aziz (Sehad & Aziz, 2016) obtained resistivity data that represents the type of subsurface rock shallow aquifer layer estimated to be composed of fine-grained sand ($0.85 \Omega\text{m}$) and sandy clay ($13.25 \Omega\text{m}$) with a depth between 7.35– 29.44 m.

From various studies that have been carried out, information about aquifer layers in Riau province is still very limited, aquifer mapping is only carried out in the area around city of Pekanbaru (Alfadli et al., 2017; Parhusip & Syech, 2013), while information on the location of aquifers is also needed in other districts in Riau Province, one of them is Rokan Hulu District. This district is one of the twelve districts in the city in Riau Province. Some clean water facilities in this district do not operate optimally because the source of water used is based on wells. Water from

the wellbore is no longer flowing because the determination of the drilling location is only an estimate and is not well planned. With this problem, information about the area of the aquifer is needed.

Masda Makmur village is one of the communities located in Rambah Samo District, Rokan Hulu Regency. This village is located not far from the river, but during the dry season, the village often experiences drought. Most of the residents use rain-fed wells as water reserves for their daily needs. This is indeed not effective, because, in the end, these wells will experience drought during the dry season. To be able to minimize dryness during the dry season, local people can use groundwater sources. But until now, no research has been carried out to detect the presence of groundwater in this area.

METHODS

The research was conducted in Masda Makmur Village, Rokan Hulu Regency. The tool used in this research is resistivity meter Georesist RS505, GPS, and Laptop. The tools used in processing data are Progress and IP2WIN software.

The geological map of the research area is shown in Figures 1 and 2. The geological setting is divided into many stratigraphic components (Sihapas Formation Formation, Farmer Formation, Alluvium, and Minas Formation). The hydrological characteristics of Rokan Hulu are shown by several rivers flowing from the peak of Suligi or from the east to the hills towards the West.

The research area was around the river. The river flowing in Rokan Hulu consists of two types of streams, the river flowing through Rokan Hulu and the river flowing only in Rokan Hulu alone. The method used in this study is the resistivity geoelectric method using the Schlumberger configuration.




Figure 1. Geological Map of the Research Area




Figure 2. Map Based on The Points Electrodes

The resistivity geoelectric method is still the most powerful technique and cost-effective in the investigation of groundwater (Mahmoud & Kotb, 2017). All types of resistivity measurements are seeking to track changes in resistivity values vertically

and horizontally (Bashir et al., 2014; Kumar, 2015; Mahmoud & Kotb, 2017).

This method consists of electric current between two electrodes injected through the ground, and then induction between two potential electrodes produces a potential difference (Claude, Théophile, Patrick, &

Crepin, 2014; Tijani, Osinowo, & Ogedengbe, 2009). The geoelectric data collection location map based on the points of the electrodes which are formed which form several lines (Figure 2. Map of Research).

The electrode arrangement in the Schlumberger configuration can be seen in Figure 3. The type of material can be known based on the resistivity value of the structure in each layer presented in table 1.

From Figure 3 above, the apparent resistivity equation for this configuration is formulated as (Fitrianto, Taufiq, & Mukromin, 2018):

$$\rho_a = K \frac{\Delta V}{I} \quad (1)$$

With ρ_a is pseudo resistivity (Ωm), K is a geometry factor, ΔV is a potential difference (Volt), and I is the amount of current (ampere).




Figure 3. Arrangement of Electrodes in Schlumberger Configuration

Table 1. The Resistivity of Material (Febriani and Sohibun, 2019)

Material	Resistivity (Ωm)
Igneous and Metamorphic	
Rocks	$5 \times 10^3 - 10^6$
Granite	$10^3 - 10^6$
Basalt	$6 \times 10^2 - 4 \times 10^7$
Slate	$10^2 - 2,5 \times 10^8$
Marble	$10^2 - 2 \times 10^8$
Quartzite	$50 - 4 \times 10^2$
Sedimentary Rocks	
Sandstone	$8 - 4 \times 10^3$
Shale	$20 - 2 \times 10^3$
Limestone	$50 - 4 \times 10^2$
Soils and water	
Clay	$1 - 100$
Alluvium	$10 - 800$
Groundwater	$10 - 100$
Seawater	$0,2$

RESULTS AND DISCUSSION

Sounding 1

On sounding 1, the track length is 100 meters. This track is located at 0054' 34'E and 100023'44' N. The interpretation results on track one is shown in Figure 4 and 7(a) with an RMS error of 9,08%.

In the first layer with a depth of 0-0,5 m, the resistivity value is 79,90 Ωm . This type of resistivity value is interpreted as a surface groundwater layer which is thought to originate from rainwater infiltration (Bakri, 2019). At a depth of 0,5-3 m has a resistivity value of 154,69 Ωm . This type of resistivity value is associated with alluvium. The kind of resistivity value is above 100 Ωm .

The resistance value of this type is higher than the value of the previous resistance (Febriani & Sohibun, 2019). Presumably, this layer is the gravel layer. At depths of 3 – 8,5 m have a more significant value of resistivity of 1535,27 Ωm . This high type of resistivity value is assumed to be a sandstone layer. This layer does not contain groundwater. At a depth of 8,5 - 15 m has a resistivity value of 154,91 Ωm . This type of resistance value is thought to be a layer of gravel (Bakri, 2019) and at a depth of 15-22 m has a low resistance value of 7,44 Ωm .

The value of this type of low resistivity is associated with groundwater, namely (1-100 Ωm). According to Wijaya (Wijaya, 2015), the value of detainees of types less than (10 Ωm) is assumed to be groundwater that has poor quality. Seen conditions in the field, there is a pile of garbage that is thought to reduce the quality of the groundwater.

Sounding 2

On sounding 2, the track length is 100 meters. This track is located at coordinates 0054' 37" E and 100023'43" N. The results of the interpretation on the second track with RMS error 7,65% have different layer variations shown in Figures 5 and 7 (b). In the first layer, with a depth of 0 – 0,5 m has a resistivity value of 20,59 Ωm .

This type of resistivity value is interpreted as a surface groundwater layer which is thought to originate from rainwater infiltration (Bakri, 2019). At a depth of 0,5 - 3 m has a resistance value of 449,86 Ωm . Alleged rock in this layer is gravel (Febriani & Sohibun, 2019). At a depth of 3,5-13 m has a higher resistance value than the previous layer, which is 2874,68 Ωm . This type of resistivity value can be interpreted as a layer of sandstone that has no potential for groundwater.

At a depth of approximately 13-18 m, the value of the resistivity is lower, which is 75,73 Ωm . This layer is supposed to be a layer of sand that can escape and store water. This type of resistivity value is assumed to be potential groundwater because at this depth the layer thickness is up to 5 meters (Febriani & Sohibun, 2019).

Sounding 3

The length of the third tracks stretches with a length of 100 meters with a distance between electrodes of 0,5 meters. This track is located in coordinates 0054'36" E, and 100023'40" N. The interpretation of the three tracks with 4,3% RMS errors have different layer variations (Figures 6 and 7 (c)).

In the first, second, and third layers, it has a resistivity value of 183 Ωm , 377,27 Ωm , and 687,34 Ωm . These three layers are thought to be non-aqueous layers because they have resistivity values of more than 100 μm (Febriani & Sohibun, 2019). From a depth of 7,5-28 m, the type of resistivity value between 42-93,26 Ωm can be interpreted as one variation of the layer, namely the sand layer with a layer thickness of up to 20 m. The depth of this layer can be assumed as potential groundwater (Bakri, 2019).




Figure 4. Results of Data Processing Resistivity to Depth on Sounding 1




Figure 5. Results of Data Processing Resistivity to Depth on Sounding 2







Figure 6. Results of Data Processing Resistivity to Depth on Sounding 3



(a)



(b)



(c)

Figure 7.(a) 2D cross section sounding 1, (b) 2D cross section sounding 2, (c) 2D cross section sounding 3

CONCLUSION AND SUGGESTION

Based on the results of the research, it can be concluded that the groundwater was found in all three soundings points with varying resistivity values ranging from 7 to 93.26 Ω m. The type of constituent material is sand. The three sounding points have different layers of variation. The deeper resistivity value and the lower it shows that the layer has the potential as a carrier layer of groundwater (aquifer). Sounding 1 and 2 have the potential to make bore wells which are thought to be depressed aquifer while sounding three is believed to be a free aquifer. Information on aquifer layers that have been obtained from the research can be a reference for stakeholders and the community to map the location of drilling wells.

AUTHOR CONTRIBUTIONS

YF drafted and designed the research design. RA detects the presence of water and collects data. AA performs statistical calculations and generates results. MA conducts discussions. DN made edits.

REFERENCES

- Alfadli, M. K., Mardiana, U., Hadian, M. S. D., Mohammad, F., Natasia, N., & Imaduddin, M. (2017). Pemetaan Cekungan Airtanah Pekanbaru Menggunakan Data VES (Vertical Electrical Sounding), Provinsi Riau, Indonesia Pekanbaru Groundwater Basin Mapping Using VES (Vertical Electrical Sounding) Data, Riau Province, Indonesia. *OSF Repository*.
- Bakri, H. (2019). Pendugaan Ketebalan Aquifer Air Tanah untuk Pengembangan Kawasan Sofifi Maluku Utara. *Jurnal Geomine*, 4(1), 5-10. <https://doi.org/10.33536/jg.v4i1.37>
- Bashir, I. Y., Izham, M. Y., & Main, R. (2014). Vertical Electrical Sounding Investigation of Aquifer Composition and it's Potential to Yield Groundwater in Some Selected Towns in Bida Basin of North Central Nigeria. *Journal of Geography and Geology*, 6(1), 60–69. <https://doi.org/10.5539/jgg.v6n1p60>
- Claude, N. P., Théophile, N.-M., Patrick, A. S., & Crepin, K. T. (2014). Evidence of Iron Mineralization Channels in the Messondo Area (Centre-Cameroon) Using Geoelectrical (DC & IP) Methods: A Case Study. *International Journal of Geosciences*, 05(03), 346–361. <https://doi.org/10.4236/ijg.2014.53034>
- Devi, A., Israil, M., Anbalagan, R., & Gupta, P. K. (2017). Subsurface Soil Characterization Using Geoelectrical and Geotechnical Investigations at a Bridge Site in Uttarakhand Himalayan Region. *Journal of Applied Geophysics*, 144, 78–85. <https://doi.org/10.1016/j.jappgeo.2017.07.005>
- Edisar, M. (2013). Pemetaan Zonasi Air Bawah Tanah di Kecamatan Pinggir Kabupaten Bengkalis Provinsi Riau. *Prosiding SEMIRATA 2013*, 405–408. Retrieved from <http://jurnal.fmipa.unila.ac.id/index.php/semirata/article/view/767>
- El-Hameed, A. G. A., El-Shayeb, H. M., El-Araby, N. A., & Hegab, M. G. (2017). Integrated Geoelectrical and Hydrogeological Studies on Wadi Qena, Egypt. *NRIAG Journal of Astronomy and Geophysics*, 6(1), 218–229. <https://doi.org/10.1016/j.nrjag.2017.03.003>
- Febriani, Y., & Sohibun. (2019). Aplikasi Metode Geolistrik Konfigurasi Schlumberger untuk Mengidentifikasi Lapisan Air Tanah di Desa Ulak Patian Rokan Hulu Riau. *Jurnal Fisika Flux*, 16(1), 54-60.
- Fitrianto, T. N., Supriyadi, S., Taufiq, U. A., Mukromin, T. M., & Wardana, A. P. (2018). Identifikasi Potensi Air Tanah Menggunakan Metode Geolistrik Resistivitas Konfigurasi Schlumberger di Kelurahan Bapangsari Kecamatan Bagelen Kabupaten Purworejo. *Jurnal*

- Fisika Flux: Jurnal Ilmiah Fisika FMIPA Universitas Lambung Mangkurat*. 15 (2), 100–104.
- Hakim, H., & Manrulu, R. H. (2017). Aplikasi Konfigurasi Wenner dalam Menganalisis Jenis Material Bawah Permukaan. *Jurnal Ilmiah Pendidikan Fisika Al-Biruni*, 5(1), 95. <https://doi.org/10.24042/jpifalbiruni.v5i1.109>
- Hanifa, D., Sota, I., & Siregar, S. S. (2016). Penentuan Lapisan Akuifer Air Tanah dengan Metode Kalimantan Selatan. *Jurnal Fisika Flux: Jurnal Ilmiah Fisika FMIPA Universitas Lambung Mangkurat*. 13 (1) 30–39.
- Helaly, A. S. (2017). Assessment of Groundwater Potentiality Using Geophysical Techniques in Wadi Allaqi Basin, Eastern Desert, Egypt – Case Study. *NRIAG Journal of Astronomy and Geophysics*, 6(2), 408–421. <https://doi.org/10.1016/j.nrjag.2017.09.003>
- Heradian, E. A., & Arman, Y. (2015). Pendugaan Bidang Gelincir Tanah Longsor di Desa Aruk Kecamatan Sajingan Besar Kabupaten Sambas dengan Menggunakan Metode Tahanan Jenis. *III*(2), 56–61.
- Hewaidy, A. G. A., El-Motaal, E. A., Sultan, S. A., Ramdan, T. M., El khafif, A. A., & Soliman, S. A. (2015). Groundwater Exploration using Resistivity and Magnetic Data at the Northwestern Part of the Gulf of Suez, Egypt. *Egyptian Journal of Petroleum*, 24(3), 255–263. <https://doi.org/10.1016/j.ejpe.2015.07.010>
- Hulu, Bappeda Rokan. (2015). Program Percepatan Sanitasi Pemukiman (PPSP). *Laporan Pokja Air Minum dan Penyehatan Lingkungan*.
- Hulu, BPS Rokan. (2017). Kabupaten Rokan Hulu dalam Angka. *Laporan BPS Rokan Hulu*, 14070.1702(1), 228.
- Jusuf, A. S. (2015). Geologi dan Pemetaan Cekungan Air Tanah Kota Gorontalo dengan Menggunakan Metode Geolistrik Tahanan Jenis. *Jurnal Fakultas MIPA UNG*.
- Kumar, R. (2015). Geohydrological Investigation Using Vertical Electrical Sounding at Banaras Hindu University Campus, Varanasi, U.P, India. *International Journal of Research in Engineering and Technology*, 03(09), 252–256. <https://doi.org/10.15623/ijret.2014.0309038>
- Mahmoud, H. H., & Kotb, A. D. M. (2017). Impact of the Geological Structures on the Groundwater potential Using eophysical Techniques in West Bani Mazar area, El Minia – Western Desert, Egypt. *Journal of African Earth Sciences*, 130, 161–173. <https://doi.org/10.1016/j.jafrearsci.2017.03.024>
- Mohamaden, M. I. I. (2016). Delineating Groundwater Aquifer and Subsurface Structures by Using Geoelectrical Data: Case Study (Dakhla Oasis, Egypt). *NRIAG Journal of Astronomy and Geophysics*, 5(1), 247–253. <https://doi.org/10.1016/j.nrjag.2016.05.001>
- Parhusip, M., & Syech, R. (2013). Menentukan Akuifer Lapisan Air Tanah dengan Metode Geolistrik Konfigurasi Schlumberger di Perumahan Griyo Puspito dan Bumi Tampan Lestari. *FMIPA Universitas Riau*, 1–8.
- Saranga, H. T., As'ari, & Tongkukut, S. H. J. (2016). Deteksi Air Tanah Menggunakan Metode Geolistrik Resistivitas Konfigurasi Wenner-Schlumberger di Masjid Kampus Universitas Sam Ratulangi dan Sekitarnya. *Jurnal Mipa Unsrat*, 5(2), 70–75.
- Sedana, D., As'ari, & Tanauma, A. (2015). The Mapping of Groundwater Aquifers at the Ringroad Malendeng Village by Using Geoelectric Resistivity Method.

- Jurnal Ilmiah Sains*, 15(2), 1–5.
- Sehah, S., & Aziz, A. N. (2016). Pendugaan Kedalaman Air Tanah Menggunakan Metode Geolistrik Konfigurasi Schlumberger di Desa Bojongsari, Kecamatan Alian, Kabupaten Kebumen. *Jurnal Neutrino*, 8(2), 41. <https://doi.org/10.18860/neu.v8i2.3239>
- Tijani, M. N., Osinowo, O. O., & Ogedengbe, O. (2009). Mapping of Sub-Surface Fracture Systems Using Integrated Electrical Resistivity Profiling and VLF-EM Methods: A Case Study of Suspected Gold Mineralization. *RMZ–Materials and Geoenvironment*, 56(4), 415–436.
- Wijaya, A. S. (2015). Aplikasi Metode Geolistrik Resistivitas Konfigurasi Wenner untuk Menentukan Struktur Tanah di Halaman Belakang SCC ITS Surabaya. *Jurnal Fisika Indonesia*, 19(55), 1–5.

Determination of Groundwater Using Geoelectric Methods

by M Apriniyadi

Submission date: 29-Mar-2023 12:47PM (UTC+0700)

Submission ID: 2049759328

File name: Artikel_Albiruni_Rokan_Hulu.pdf (836.74K)

Word count: 3301

Character count: 17463



Determination of Groundwater Using Geoelectric Methods: Schlumberger Configuration in Rokan Hulu Regency

Yeza Febriani^{*1}, Riza Ari Rohman², Azmi Azra³, Mohammad Apriniyadi⁴, Donata Nariswari Wahyu Wardani⁵

^{1,2,3}Physics Study Programme, Faculty of teacher training and education, Pasir Pengaraian University, Rokan Hulu Regency 28557, Riau Province, Indonesia;

⁴Geology Technic Programme, Universitas Trisakti, Jakarta Barat, Jakarta, Indonesia;

⁵Groundwater management Programme, Faculty of Geoscience, Geoengineering and Mining, Freiberg University of Mining and Technology, Akademiestraße 609599 Freiberg, Germany.

*Corresponding Address: yezafebriani@upp.ac.id

Article Info

Article history:

Received: April 1st, 2019

Accepted: October 20th, 2019

Published: October 30th, 2019

Keywords:

aquifer;
geoelectric;
groundwater;
resistivity;
schlumberger.

ABSTRACT

This study aimed at detecting the presence of groundwater in Masda Makmur Village, Rambah Samo District, Rokan Hulu Regency, using the Schlumberger configuration geoelectric method. The study consisted of 3 sounding points within every 100 m sounding track length. The measurement data is processed using IP2WIN software and Progress to see layer data below the ground level based on the value of the resistivity type. The results showed that the estimated groundwater could be found on track one with a kind of resistivity value of 7,44 Ω m at a depth of 15-22 m. Alleged groundwater can be found on the second track with a type of resistivity value of 75,73 Ω mat a depth of 13-18 m, and the estimated groundwater can be located on the third track with a kind of resistivity value of 82,52 Ω m, 93,26 Ω m. The deeper and the lower resistivity value shows that the layer has the potential as a carrier layer of groundwater (aquifer). Sounding 1 and 2 have the potential to make bore wells which are thought to be depressed aquifer. The results of this study can be used further for mapping the location of community boreholes

© 2019 Physics Education Department, UIN Raden Intan Lampung, Indonesia.

INTRODUCTION

Water is essential for life on earth (Saranga et al., 2016). Water consists of two atoms of hydrogen and one atom of oxygen (H₂O). Living things cannot live without water. The increasing population requires a sufficient amount of water. In Indonesia, especially Rokan Hulu Regency, population growth is increasing every year (Bappeda Rokan Hulu, 2015).

In 2016, the percentage of population growth rate in the Rokan Hulu Regency was 0.27%, 2017 experienced an increase of 0.09%, and in 2018 there was an increase of

0.05%. For Rambah Samo District, Rokan Hulu Regency in 2016 the population growth rate was 1,36%, 2017 increased by 1,82% and in 2018 again experienced a significant increase of 2,05% (BPS Rokan Hulu, 2017). The higher of the population growth rate, the more need for clean water by the community. Groundwater is one source of life necessities for living things on earth (Hanifa, Sota, & Siregar, 2016; Sehad & Aziz, 2016). Groundwater is all water found in layers containing water (aquifers) below the surface of the soil. The role of groundwater is increasingly essential to

How to cite

Febriani, Y., Rohman, R. A., Asra, A., Apriniyadi, M., & Wardani, D. N. W. (2019). Determination of Groundwater Using Geoelectric Methods: Schlumberger Configuration in Rokan Hulu Regency. *Jurnal Ilmiah Pendidikan Fisika Al-Biruni*, 8(2), 141-152.

become the primary water source to fulfill the basic needs of many people (Jusuf, 2015; Sedana, As'ari, & Tanaur, 2015).

Specific methods can detect the presence of groundwater in the subsurface layer (Devi, Israil, Anbalagan, & Gupta, 2017; Heradian & Arman, 2015; Wijaya, 2015). One of which is by using the geoelectric resistivity method. This method is the most popular method of investigating groundwater from the surface of the earth (Bakri, 2019; Bashir, Izham, & Main, 2014; El-Hameed, El-Shayeb, El-Araby, & Hegab, 2017; Helaly, 2017; Hewaidy et al., 2015; Jusuf, 2015; Mohamaden, 2016). The geoelectric method is a method that studies the nature of electricity on the planet by detecting it on the earth's surface (Hakim & Manrulu, 2017).

Several studies on the detection of groundwater estimation using Schlumberger configuration have been carried out, including editions (Edisar, 2013) of data obtained, namely zoning of free aquifers and depressed aquifers. Sehad and Aziz (Sehad & Aziz, 2016) obtained resistivity data that represents the type of subsurface rock shallow aquifer layer estimated to be composed of fine-grained sand ($0.85 \Omega\text{m}$) and sandy clay ($13.25 \Omega\text{m}$) with a depth between 7.35– 29.44 m.

From various studies that have been carried out, information about aquifer layers in Riau province is still very limited, aquifer mapping is only carried out in the area around city of Pekanbaru (Alfadli et al., 2017; Parhusip & Syech, 2013), while information on the location of aquifers is also needed in other districts in Riau Province, one of them is Rokan Hulu District. This district is one of the twelve districts in the city in Riau Province. Some clean water facilities in this district do not operate optimally because the source of water used is based on wells. Water from

the wellbore is no longer flowing because the determination of the drilling location is only an estimate and is not well planned. With this problem, information about the area of the aquifer is needed.


Masda Makmur village is one of the communities located in Rambah Samo District, Rokan Hulu Regency. This village is located not far from the river, but during the dry season, the village often experiences drought. Most of the residents use rain-fed wells as water reserves for their daily needs. This is indeed not effective, because, in the end, these wells will experience drought during the dry season. To be able to minimize dryness during the dry season, local people can use groundwater sources. But until now, no research has been carried out to detect the presence of groundwater in this area.

METHODS

The research was conducted in Masda Makmur Village, Rokan Hulu Regency. The tool used in this research is resistivity meter Georesist RS505, GPS, and Laptop. The tools used in processing data are Progress and P2WIN software.

The geological map of the research area is shown in Figures 1 and 2. The geological setting is divided into many stratigraphic components (Sihapas Formation Formation, Farmer Formation, Alluvium, and Minas Formation). The hydrological characteristics of Rokan Hulu are shown by several rivers flowing from the peak of Suligi or from the east to the hills towards the West.

The research area was around the river. The river flowing in Rokan Hulu consists of two types of streams, the river flowing through Rokan Hulu and the river flowing only in Rokan Hulu alone. The method used in this study is the resistivity geoelectric method using the Schlumberger configuration.



4
Figure 1. Geological Map of the Research Area




Figure 2. Map Based on The Points Electrodes

The resistivity geoelectric method is still the most powerful technique and cost-effective in the investigation of groundwater (Mahmoud & Kotb, 2017). All types of resistivity measurements are seeking to track changes in resistivity values vertically

and horizontally (Bashir et al., 2014; Kumar, 2015; Mahmoud & Kotb, 2017).

This method consists of electric current between two electrodes injected through the ground, and then induction between two potential electrodes produces a potential difference (Claude, Théophile, Patrick, &

Crepin, 2014; Tijani, Osinowo, & Ogedengbe, 2009). The geoelectric data collection location map based on the points of the electrodes which are formed which form several lines (Figure 2. Map of Research).

The electrode arrangement in the Schlumberger configuration can be seen in Figure 3. The type of material can be known based on the resistivity value of the structure in each layer presented in table 1.

From Figure 3 above, the apparent resistivity equation for this configuration is formulated as (Fitrianto, Taufiq, & Mukromin, 2018):

$$\rho_a = K \frac{\Delta V}{I} \tag{1}$$

With ρ_a is pseudo resistivity (Ωm), K is a geometry factor, ΔV is a potential difference (Volt), and I is the amount of current (ampere).




Figure 3. Arrangement of Electrodes in Schlumberger Configuration

Table 1. The Resistivity of Material (Febriani and Sohibun, 2019)

Material	Resistivity (Ωm)
Igneous and Metamorphic	
Rocks	$5 \times 10^3 - 10^6$
Granite	$10^3 - 10^6$
Basalt	$6 \times 10^2 - 4 \times 10^7$
Slate	$10^2 - 2,5 \times 10^8$
Marble	$10^2 - 2 \times 10^8$
Quartzite	$50 - 4 \times 10^2$
Sedimentary Rocks	
Sandstone	$8 - 4 \times 10^3$
Shale	$20 - 2 \times 10^3$
Limestone	$50 - 4 \times 10^2$
Soils and water	
Clay	1 - 100
Alluvium	10 - 800
Groundwater	10-100
Seawater	0,2

RESULTS AND DISCUSSION

Sounding 1

On sounding 1, the track length is 100 meters. This track is located at 0054' 34'E and 100023'44' N. The interpretation results on track one is shown in Figure 4 and 7(a) with an RMS error of 9,08%.

In the first layer with a depth of 0-0,5 m, the resistivity value is 79,90 Ωm . This type of resistivity value is interpreted as a surface groundwater layer which is thought to originate from rainwater infiltration (Bakri, 2019). At a depth of 0,5-3 m has a resistivity value of 154,69 Ωm . This type of resistivity value is associated with alluvium. The kind of resistivity value is above 100 Ωm .

The resistance value of this type is higher than the value of the previous resistance (Febriani & Sohibun, 2019). Presumably, this layer is the gravel layer. At depths of 3 - 8,5 m have a more significant value of resistivity of 1535,27 Ωm . This high type of resistivity value is assumed to be a sandstone layer. This layer does not contain groundwater. At a depth of 8,5 - 15 m has a resistivity value of 154,91 Ωm . This type of resistance value is thought to be a layer of gravel (Bakri, 2019) and at a depth of 15-22 m has a low resistance value of 7,44 Ωm .

The value of this type of low resistivity is associated with groundwater, namely (1-100 Ωm). According to Wijaya (Wijaya, 2015), the value of detainees of types less than (10 Ωm) is assumed to be groundwater that has poor quality. Seen conditions in the field, there is a pile of garbage that is thought to reduce the quality of the groundwater.

Sounding 2

On sounding 2, the track length is 100 meters. This track is located at coordinates 0054' 37" E and 100023'43" N. The results of the interpretation on the second track with RMS error 7,65% have different layer variations shown in Figures 5 and 7 (b) in the first layer, with a depth of 0 - 0,5 m has a resistivity value of 20,59 Ωm .

This type of resistivity value is interpreted as a surface groundwater layer which is thought to originate from rainwater infiltration (Bakri, 2019). At a depth of 0,5 - 3 m has a resistance value of 449,86 Ω m. Alleged rock in this layer is gravel (Febriani & Sohibun, 2019). At a depth of 3,5-13 m has a higher resistance value than the previous layer, which is 2874,68 Ω m. This type of resistivity value can be interpreted as a layer of sandstone that has no potential for groundwater.

At a depth of approximately 13-18 m, the value of the resistivity is lower, which is 75,73 Ω m. This layer is supposed to be a layer of sand that can escape and store water. This type of resistivity value is assumed to be potential groundwater because at this depth the layer thickness is up to 5 meters (Febriani & Sohibun, 2019).

Sounding 3

The length of the third tracks stretches with a length of 100 meters with a distance between electrodes of 0,5 meters. This track is located in coordinates 0054'36" E, and 100023'40" N. The interpretation of the three tracks with 4,3% RMS errors have different layer variations (Figures 6 and 7 (c)).

In the first, second, and third layers, it has a resistivity value of 183 Ω m, 377,27 Ω m, and 687,34 Ω m. These three layers are thought to be non-aqueous layers because they have resistivity values of more than 100 μ m (Febriani & Sohibun, 2019). From a depth of 7,5-28 m, the type of resistivity value between 42-93,26 Ω m can be interpreted as one variation of the layer, namely the sand layer with a layer thickness of up to 20 m. The depth of this layer can be assumed as potential groundwater (Bakri, 2019).




Figure 4. Results of Data Processing Resistivity to Depth on Sounding 1




Figure 5. Results of Data Processing Resistivity to Depth on Sounding 2







Figure 6. Results of Data Processing Resistivity to Depth on Sounding 3



(a)



(b)



(c)

Figure 7.(a) 2D cross section sounding 1, (b) 2D cross section sounding 2, (c) 2D cross section sounding 3

12

CONCLUSION AND SUGGESTION

Based on the results of the research, it can be concluded that the groundwater was found in all three soundings points with varying resistivity values ranging from 7 to 93.26 Ω m. The type of constituent material is sand. The three sounding points have different layers of variation. The deeper resistivity value and the lower it shows that the layer has the potential as a carrier layer of groundwater (aquifer). Sounding 1 and 2 have the potential to make bore wells which are thought to be depressed aquifer while sounding three is believed to be a free aquifer. Information on aquifer layers that have been obtained from the research can be a reference for stakeholders and the community to map the location of drilling wells.

AUTHOR CONTRIBUTIONS

YF drafted and designed the research design. RA detects the presence of water and collects data. AA performs statistical calculations and generates results. MA conducts discussions. DN made edits.

REFERENCES

- Alfadli, M. K., Mardiana, U., Hadian, M. S. D., Mohammad, F., Natasia, N., & Imaduddin, M. (2017). Pemetaan Cekungan Airtanah Pekanbaru Menggunakan Data VES (Vertical Electrical Sounding), Provinsi Riau, Indonesia Pekanbaru Groundwater Basin Mapping Using VES (Vertical Electrical Sounding) Data, Riau Province, Indonesia. *OSF Repository*.
- Bakri, H. (2019). Pendugaan Ketebalan Aquifer Air Tanah untuk Pengembangan Kawasan Sofifi Maluku Utara. *Jurnal Geomine*, 4(1), 5-10. <https://doi.org/10.33536/jg.v4i1.37>
- Bashir, I. Y., Izham, M. Y., & Main, R. (2014). Vertical Electrical Sounding Investigation of Aquifer Composition and it's Potential to Yield Groundwater in Some Selected Towns in Bida Basin of North Central Nigeria. *Journal of Geography and Geology*, 6(1), 60–69. <https://doi.org/10.5539/jgg.v6n1p60>
- Claude, N. P., Théophile, N.-M., Patrick, A. S., & Crepin, K. T. (2014). Evidence of Iron Mineralization Channels in the Messondo Area (Centre-Cameroon) Using Geoelectrical (DC & IP) Methods: A Case Study. *International Journal of Geosciences*, 05(03), 346–361. <https://doi.org/10.4236/ijg.2014.53034>
- Devi, A., Israil, M., Anbalagan, R., & Gupta, P. K. (2017). Subsurface Soil Characterization Using Geoelectrical and Geotechnical Investigations at a Bridge Site in Uttarakhand Himalayan Region. *Journal of Applied Geophysics*, 144, 78–85. <https://doi.org/10.1016/j.jappgeo.2017.07.005>
- Edisar, M. (2013). Pemetaan Zonasi Air Bawah Tanah di Kecamatan Pinggir Kabupaten Bengkalis Provinsi Riau. *Prosiding SEMIRATA 2013*, 405–408. Retrieved from <http://jurnal.fmipa.unila.ac.id/index.php/semirata/article/view/767>
- El-Hameed, A. G. A., El-Shayeb, H. M., El-Araby, N. A., & Hegab, M. G. (2017). Integrated Geoelectrical and Hydrogeological Studies on Wadi Qena, Egypt. *NRIAG Journal of Astronomy and Geophysics*, 6(1), 218–229. <https://doi.org/10.1016/j.nrjag.2017.03.003>
- Febriani, Y., & Sohibun. (2019). Aplikasi Metode Geolistrik Konfigurasi Schlumberger untuk Mengidentifikasi Lapisan Air Tanah di Desa Ulak Patian Rokan Hulu Riau. *Jurnal Fisika Flux*, 16(1), 54-60.
- Fitrianto, T. N., Supriyadi, S., Taufiq, U. A., Mukromin, T. M., & Wardana, A. P. (2018). Identifikasi Potensi Air Tanah Menggunakan Metode Geolistrik Resistivitas Konfigurasi Schlumberger di Kelurahan Bapangsari Kecamatan Bagelen Kabupaten Purworejo. *Jurnal*

- Fisika Flux: Jurnal Ilmiah Fisika FMIPA Universitas Lambung Mangkurat*. 15 (2), 100–104.
- Hakim, H., & Manrulu, R. H. (2017). Aplikasi Konfigurasi Wenner dalam Menganalisis Jenis Material Bawah Permukaan. *Jurnal Ilmiah Pendidikan Fisika Al-Biruni*, 5(1), 95. <https://doi.org/10.24042/jpifalbiruni.v5i1.109>
- Hanifa, D., Sota, I., & Siregar, S. S. (2016). Penentuan Lapisan Akuifer Air Tanah dengan Metode Kalimantan Selatan. *Jurnal Fisika Flux: Jurnal Ilmiah Fisika FMIPA Universitas Lambung Mangkurat*. 13 (1) 30–39.
- Helaly, A. S. (2017). Assessment of Groundwater Potentiality Using Geophysical Techniques in Wadi Allaqi Basin, Eastern Desert, Egypt – Case Study. *NRIAG Journal of Astronomy and Geophysics*, 6(2), 408–421. <https://doi.org/10.1016/j.nrjag.2017.09.003>
- Heradian, E. A., & Arman, Y. (2015). Pendugaan Bidang Gelincir Tanah Longsor di Desa Aruk Kecamatan Sajingan Besar Kabupaten Sambas dengan Menggunakan Metode Tahanan Jenis. *III*(2), 56–61.
- Hewaify, A. G. A., El-Motaal, E. A., Sultan, S. A., Ramdan, T. M., El khafif, A. A., & Soliman, S. A. (2015). Groundwater Exploration using Resistivity and Magnetic Data at the Northwestern Part of the Gulf of Suez, Egypt. *Egyptian Journal of Petroleum*, 24(3), 255–263. <https://doi.org/10.1016/j.ejpe.2015.07.010>
- Hulu, Bappeda Rokan. (2015). Program Percepatan Sanitasi Pemukiman (PPSP). *Laporan Pokja Air Minum dan Penyehatan Lingkungan*.
- Hulu, BPS Rokan. (2017). Kabupaten Rokan Hulu dalam Angka. *Laporan BPS Rokan Hulu*, 14070.1702(1), 228.
- Jusuf, A. S. (2015). Geologi dan Pemetaan Cekungan Air Tanah Kota Gorontalo dengan Menggunakan Metode Geolistrik Tahanan Jenis. *Jurnal Fakultas MIPA UNG*.
- Kumar, R. (2015). Geohydrological Investigation Using Vertical Electrical Sounding at Banaras Hindu University Campus, Varanasi, U.P, India. *International Journal of Research in Engineering and Technology*, 03(09), 252–256. <https://doi.org/10.15623/ijret.2014.0309038>
- Mahmoud, H. H., & Kotb, A. D. M. (2017). Impact of the Geological Structures on the Groundwater potential Using eophysical Techniques in West Bani Mazar area, El Minia – Western Desert, Egypt. *Journal of African Earth Sciences*, 130, 161–173. <https://doi.org/10.1016/j.jafrearsci.2017.03.024>
- Mohamaden, M. I. I. (2016). Delineating Groundwater Aquifer and Subsurface Structures by Using Geoelectrical Data: Case Study (Dakhla Oasis, Egypt). *NRIAG Journal of Astronomy and Geophysics*, 5(1), 247–253. <https://doi.org/10.1016/j.nrjag.2016.05.001>
- Parhusip, M., & Syech, R. (2013). Menentukan Akuifer Lapisan Air Tanah dengan Metode Geolistrik Konfigurasi Schlumberger di Perumahan Griyo Puspito dan Bumi Tampan Lestari. *FMIPA Universitas Riau*, 1–8.
- Saranga, H. T., As'ari, & Tongkukut, S. H. J. (2016). Deteksi Air Tanah Menggunakan Metode Geolistrik Resistivitas Konfigurasi Wenner-Schlumberger di Masjid Kampus Universitas Sam Ratulangi dan Sekitarnya. *Jurnal Mipa Unsrat*, 5(2), 70–75.
- Sedana, D., As'ari, & Tanauma, A. (2015). The Mapping of Groundwater Aquifers at the Ringroad Malendeng Village by Using Geoelectric Resistivity Method.

- Jurnal Ilmiah Sains*, 15(2), 1–5.
- Sehah, S., & Aziz, A. N. (2016). Pendugaan Kedalaman Air Tanah Menggunakan Metode Geolistrik Konfigurasi Schlumberger di Desa Bojongsari, Kecamatan Alian, Kabupaten Kebumen. *Jurnal Neutrino*, 8(2), 41. <https://doi.org/10.18860/neu.v8i2.3239>
- Tijani, M. N., Osinowo, O. O., & Ogedengbe, O. (2009). Mapping of Sub-Surface Fracture Systems Using Integrated Electrical Resistivity Profiling and VLF-EM Methods: A Case Study of Suspected Gold Mineralization. *RMZ–Materials and Geoenvironment*, 56(4), 415–436.
- Wijaya, A. S. (2015). Aplikasi Metode Geolistrik Resistivitas Konfigurasi Wenner untuk Menentukan Struktur Tanah di Halaman Belakang SCC ITS Surabaya. *Jurnal Fisika Indonesia*, 19(55), 1–5.

Determination of Groundwater Using Geoelectric Methods

ORIGINALITY REPORT

14%

SIMILARITY INDEX

10%

INTERNET SOURCES

11%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1

pdfs.semanticscholar.org

Internet Source

3%

2

bestjournal.untad.ac.id

Internet Source

2%

3

Iin Amalia Khusna, Sorja Koesuma, Darsono. "Determination of Groundwater Potential in Pulutan Wetan Village Wuryantoro District Wonogiri Using Vertical Electrical Sounding Method", Journal of Physics: Conference Series, 2021

Publication

2%

4

journal2.um.ac.id

Internet Source

1%

5

Sandra Sandra, Sunaldi Sunaldi, Mohammad Dahlan Th. Musa, Abdullah Abdullah, Muh. Rusydi, Maskur Maskur. "Identification of Weak Zones in the Area of Sigi Regency Office Using Geoelectric Methods of Resistivity", Gravitasi, 2022

Publication

1%



6	Submitted to Universitas Jenderal Soedirman Student Paper	1 %
7	Hussein Hosni Mahmoud, Adel Diab Mohammed Kotb. "Impact of the geological structures on the groundwater potential using geophysical techniques in West Bani Mazar area, El Minia – Western Desert, Egypt", Journal of African Earth Sciences, 2017 Publication	1 %
8	ppjp.ulm.ac.id Internet Source	1 %
9	media.neliti.com Internet Source	1 %
10	G Yuliyanto, M I Nurwidyanto. "Integrated survey to identify potential groundwater aquifers in Jabungan Semarang using geoelectric and microtremor methods", Journal of Physics: Conference Series, 2021 Publication	1 %
11	Submitted to Universitas Riau Student Paper	1 %
12	journal.unhas.ac.id Internet Source	1 %

Exclude quotes Off

Exclude matches < 15 words

Exclude bibliography On

Determination of Groundwater Using Geoelectric Methods

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12
